Note

(a) Consider two gas containers at the same temperature having the same internal energy, one is kept at rest on the ground and the other is kept in a moving train. Even though the gas container in the train is moving with the speed of the train, the internal energy of the gas in it will not increase.

(b) Consider two gas containers at the same temperature having the same internal energy, one is kept on the ground and the other is kept at some height h. Even though the container at height h is having higher gravitational potential energy, this has no influence on internal energy of the gas molecules.


EXAMPLE 8.10

When you mix a tumbler of hot water with one bucket of normal water, what will be the direction of heat flow? Justify.

The water in the tumbler is at a higher temperature than the bucket of normal water. But the bucket of normal water has larger internal energy than the hot water in the tumbler. This is because the internal energy is an extensive variable and it depends on the size or mass of the system.

Even though the bucket of normal water has larger internal energy than the tumbler of hot water, heat will flow from water in the tumbler to the water in the bucket. This is because heat flows from a body at higher temperature to the one at lower temperature and is independent of internal energy of the system.

Once the heat is transferred to an object it becomes internal energy of the object. The right way to say is 'object has certain amount of internal energy'. Heat is one of the ways to increase the internal energy of a system as shown in the Figure.



Note

It is to be noted that heat does not always increases the internal energy. Later we shall see that in ideal gases during isothermal process the internal energy will not increase even though heat flows in to the system.

8.6.1 Joule's mechanical equivalent of heat

The temperature of an object can be increased by heating it or by doing some work on it. In the eighteenth century, James Prescott Joule showed that mechanical energy can be converted into internal energy and vice versa. In his experiment, two masses were attached with a rope and a paddle wheel as shown in Figure 8.19. When these masses fall through a distance h due to gravity, both the masses lose potential energy equal to 2mgh. When the masses fall, the paddle wheel turns. Due to the turning of wheel inside water, frictional force comes in between the water and the paddle wheel. This causes

UNIT-8(XI-Physics_Vol-2).indd 117 02-01-2023 15:25:45