If Q is the heat supplied to mole of a gas at constant volume and if the temperature changes by an amount ΔT , we have

$$Q = \mu C_v \Delta T. \quad (8.18)$$

By applying the first law of thermodynamics for this constant volume process (W=0, since dV=0), we have

$$Q = \Delta U - 0 \quad (8.19)$$

By comparing the equations (8.18) and (8.19),

$$\Delta U = \mu C_v \Delta T \text{ or } C_v = \frac{1}{\mu}\frac{\Delta U}{\Delta T}$$

If the limit ΔT goes to zero, we can write

$$C_v = \frac{1}{\mu}\frac{dU}{dT} \quad (8.20)$$

Since the temperature and internal energy are state variables, the above relation holds true for any process.

8.7.2 Meyer's relation

Consider μ mole of an ideal gas in a container with volume V, pressure P and temperature T.

When the gas is heated at constant volume the temperature increases by dT. As no work is done by the gas, the heat that flows into the system will increase only the internal energy. Let the change in internal energy be dU.

If C_v is the molar specific heat capacity at constant volume, from equation (8.20)

$$dU = \mu C_v dT \quad (8.21)$$

Suppose the gas is heated at constant pressure so that the temperature increases by dT. If 'Q' is the heat supplied in this process and 'dV' the change in volume of the gas.

$$Q = \mu C_p dT \quad (8.22)$$

If W is the workdone by the gas in this process, then

$$W = PdV \quad (8.23)$$

But from the first law of thermodynamics,

$$Q = dU + W \quad (8.24)$$

Substituting equations (8.21), (8.22) and (8.23) in (8.24), we get,

$$\mu C_p dT = \mu C_v dT + PdV \quad (8.25)$$

For mole of ideal gas, the equation of state is given by

$$PV = \mu RT \Rightarrow PdV + VdP = \mu RdT \quad (8.26)$$

Since the pressure is constant, dP=0

$$\therefore C_p dT = C_v dT + RdT$$

$$\therefore C_P = C_v + R \quad \text{(or)} \quad C_p - C_v = R \quad (8.27)$$

This relation is called Meyer's relation

It implies that the molar specific heat capacity of an ideal gas at constant pressure is greater than molar specific heat capacity at constant volume.

The relation shows that specific heat at constant pressure (s_p) is always greater that specific heat at constant volume (s_v).

8.8 THERMODYNAMIC PROCESSES

8.8.1 Isothermal process

It is a process in which the temperature remains constant but the pressure and volume of a thermodynamic system will change. The ideal gas equation is

UNIT-8(XI-Physics_Vol-2).indd 124 02-01-2023 15:25:50