The x-component of momentum of the molecule after collision = $-mv_x$

The change in momentum of the molecule in x direction

=Final momentum – initial momentum = $-mv_x - mv_x = -2mv_x$

According to law of conservation of linear momentum, the change in momentum of the wall = $2mv_x$

Note

In x direction, the total momentum of the system before collision is equal to momentum of the molecule (mv_x) since the momentum of the wall is zero. According to the law of conservation of momentum the total momentum of system after the collision must be equal to total momentum of system before collision. The momentum of the molecule (in x direction) after the collision is $-mv_x$ and the momentum of the wall after the collision is $2mv_x$. So total momentum of the system after the collision is $(2mv_x - mv_x) = mv_x$ which is same as the total momentum of the system before collision.

The number of molecules hitting the right side wall in a small interval of time Δt is calculated as follows.

The molecules within the distance of $v_x \Delta t$ from the right side wall and moving towards the right will hit the wall in the time interval Δt. This is shown in the Figure 9.2. The number of molecules that will hit the right side wall in a time interval Δt is equal to the product of volume ($Av_x \Delta t$) and number density of the molecules (n). Here A is area of the wall and n is number of molecules per unit volume $\left(\frac{N}{V}\right)$. We have assumed that the number density is the same throughout the cube.




Figure 9.2 Number of molecules hitting the wall

Not all the n molecules will move to the right, therefore on an average only half of the n molecules move to the right and the other half moves towards left side.

The number of molecules that hit the right side wall in a time interval Δt

$$= \frac{n}{2} A v_x \Delta t \qquad (9.1)$$

In the same interval of time Δt, the total momentum transferred by the molecules

$$\Delta p = \frac{n}{2} A v_x \Delta t \times 2mv_x = Av_x^2 mn\Delta t \qquad (9.2)$$

From Newton's second law, the change in momentum in a small interval of time gives rise to force.

The force exerted by the molecules on the wall (in magnitude)

$$F = \frac{\Delta p}{\Delta t} = nmAv_x^2 \qquad (9.3)$$

Pressure, P = force divided by the area of the wall

$$P = \frac{F}{A} = nmv_x^2 \qquad (9.4)$$

Since all the molecules are moving completely in random manner, they do not have same speed. So we can replace the term v_x^2 by the average $\overline{v_x^2}$ in equation (9.4)

UNIT-9(XI-Physics_Vol-2).indd 166 26-03-2022 17:46:45