
Note

Conservation of energy

Both the kinetic energy and potential energy are periodic functions, and repeat their values after a time period $\frac{T}{2}$. But total energy is constant for all the values of x or t. The kinetic energy and the potential energy for a simple harmonic motion are always positive. Note that kinetic energy cannot take negative value because it is proportional to the square of velocity. The measurement of any physical quantity must be a real number. Therefore, if kinetic energy is negative then the numerical value of velocity becomes an imaginary number, which is physically not acceptable. At equilibrium, it is purely kinetic energy and at extreme positions it is purely potential energy.



EXAMPLE 10.16

Compute the position of an oscillating particle when its kinetic energy and potential energy are equal.

Solution

Since the kinetic energy and potential energy of the oscillating particle are equal,

$$\frac{1}{2}m\omega^2(A^2 - x^2) = \frac{1}{2}m\omega^2 x^2$$

$$A^2 - x^2 = x^2$$

$$2x^2 = A^2$$

$$\Rightarrow x = \pm\frac{A}{\sqrt{2}}$$

10.6 TYPES OF OSCILLATIONS:

10.6.1 Free oscillations

When the oscillator is allowed to oscillate by displacing its position from equilibrium position, it oscillates with a frequency which is equal to the natural frequency of the oscillator. Such an oscillation or vibration is known as free oscillation or free vibration. In this case, the amplitude, frequency and the energy of the vibrating object remains constant.

Examples:

(i) Vibration of a tuning fork.
(ii) Vibration in a stretched string.
(iii) Oscillation of a simple pendulum.
(iv) Oscillationsof a spring-mass system.

10.6.2 Damped oscillations

During the oscillation of a simple pendulum (in previous case), we have assumed that the amplitude of the oscillation is constant and also the total energy of the oscillator is constant. But in reality, in a medium, due to the presence of friction and air drag, the amplitude of oscillation decreases as time progresses. It implies that the oscillation is not sustained and the energy of the SHM decreases gradually indicating the loss of energy. The energy lost is absorbed by the surrounding medium. This type of

UNIT-10(XI-Physics_Vol-2).indd 213 25-03-2022 20:30:31