The length of the string, L = 80 cm = 0.8 m
The mass of the string, m = 0.32 g
= 0.32×10^{-3} kg
Therefore, the linear mass density,

$$\mu = \frac{0.32\times10^{-3}}{0.8} = 0.4\times10^{-3}\, kg\, m^{-1}$$

The tension in the string, T = 80 N

$$v = \sqrt{\frac{80}{0.4\times10^{-3}}} = 447.2 \text{ m s}^{-1}$$

The wavelength corresponding to the fundamental frequency f_1 is $\lambda_1 = 2L = 2 \times 0.8 = 1.6$ m
The fundamental frequency f_1 corresponding to the wavelength λ_1

$$f_1 = \frac{v}{\lambda_1} = \frac{447.2}{1.6} = 279.5\, Hz$$

Similarly, the frequency corresponding to the second harmonics, third harmonics and fourth harmonics are

$$f_2 = 2f_1 = 559 \text{ Hz}$$

$$f_3 = 3f_1 = 838.5 \text{ Hz}$$

$$f_4 = 4f_1 = 1118 \text{ Hz}$$

11.8.5 Laws of transverse vibrations in stretched strings

There are three laws of transverse vibrations of stretched strings which are given as follows:

(i) The law of length :

For a given wire with tension T (which is fixed) and mass per unit length μ (fixed) the frequency varies inversely with the vibrating length. Therefore,

$$f \propto \frac{1}{l} \Rightarrow f = \frac{C}{l}$$

$$\Rightarrow l \times f = \text{C, where C is a constant}$$

(ii) The law of tension:

For a given vibrating length l (fixed) and mass per unit length μ (fixed) the frequency varies directly with the square root of the tension T,

$$f \propto \sqrt{T}$$

$$\Rightarrow f = A\sqrt{T}, \text{ where A is a constant}$$

(iii) The law of mass:

For a given vibrating length l (fixed) and tension T (fixed) the frequency varies inversely with the square root of the mass per unit length μ,

$$f \propto \frac{1}{\sqrt{\mu}}$$

$$\Rightarrow f = \frac{B}{\sqrt{\mu}}, \text{ where B is a constant}$$

11.9 INTENSITY AND LOUDNESS

Consider a source and two observers (listeners). The source emits sound waves which carry energy. The sound energy emitted by the source is same regardless of whoever measures it, i.e., it is independent of any observer standing in that region. But the sound received by the two observers may be different; this is due to some factors like sensitivity of ears, etc. To quantify such thing, we define two different quantities known as intensity and loudness of sound.

11.9.1 Intensity of sound

When a sound wave is emitted by a source, the energy is carried to all possible surrounding points. The average sound energy emitted or

UNIT-11(XI-Physics_Vol-2).indd 259 26-03-2022 17:25:14