$$= -a_1(b_2c_3 - b_3c_2) + b_1(a_2c_3 - a_3c_2) - c_1(a_2b_3 - a_3b_2)$$

$$= -[a_1(b_2c_3 - b_3c_2) - b_1(a_2c_3 - a_3c_2) + c_1(a_2b_3 - a_3b_2)]$$

$$= -|A|$$

Therefore, $|A_1| = -|A|$. Thus the property is verified. ■

Property 3

If there are n interchanges of rows (columns) of a matrix A then the determinant of the resulting matrix is $(-1)^n |A|$.

Property 4

If two rows (columns) of a matrix are identical, then its determinant is zero.

Verification

Let $|A| = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_2 & b_2 & c_2 \end{vmatrix}$, with 2nd and 3rd rows are identical.

Interchanging second and third rows, we get $-|A| = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_2 & b_2 & c_2 \end{vmatrix} = |A|$.

$$\Rightarrow 2|A| = 0 \Rightarrow |A| = 0.$$ ■

Property 5

If a row (column) of a matrix A is a scalar multiple of another row (or column) of A, then its determinant is zero.

Note 7.8

(i) If all entries of a row or a column are zero, then the determinant is zero.

(ii) The determinant of a triangular matrix is obtained by the product of the principal diagonal elements.

Property 6

If each element in a row (or column) of a matrix is multiplied by a scalar k, then the determinant is multiplied by the same scalar k.

Verification

Let $|A| = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$

$$= a_1(b_2c_3 - b_3c_2) - b_1(a_2c_3 - a_3c_2) + c_1(a_2b_3 - a_3b_2)$$

Let $|A_1| = \begin{vmatrix} ka_1 & kb_1 & kc_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$

$$= ka_1(b_2c_3 - b_3c_2) - kb_1(a_2c_3 - a_3c_2) + kc_1(a_2b_3 - a_3b_2) = k|A|$$

$$= k[a_1(b_2c_3 - b_3c_2) - b_1(a_2c_3 - a_3c_2) + c_1(a_2b_3 - a_3b_2)] = k|A|$$

$$\Rightarrow |A_1| = k|A|.$$ ■

07_11th-Maths-Vol2-Unit7.indd 25 16-12-2022 10:59:49