(18) If $|\vec{a}|=13, |\vec{b}|=5$ and $\vec{a}\cdot\vec{b}=60°$ then $|\vec{a}\times\vec{b}|$ is

(1) 15 (2) 35 (3) 45 (4) 25

(19) Vectors $\vec{a}$ and $\vec{b}$ are inclined at an angle $\theta=120°$. If $|\vec{a}|=1, |\vec{b}|=2$, then $[(\vec{a}+3\vec{b})\times(3\vec{a}-\vec{b})]^2$ is equal to

(1) 225 (2) 275 (3) 325 (4) 300

(20) If $\vec{a}$ and $\vec{b}$ are two vectors of magnitude 2 and inclined at an angle $60°$, then the angle between $\vec{a}$ and $\vec{a}+\vec{b}$ is

(1) 30° (2) 60° (3) 45° (4) 90°

(21) If the projection of $5\hat{i}-\hat{j}-3\hat{k}$ on the vector $\hat{i}+3\hat{j}+\lambda\hat{k}$ is same as the projection of $\hat{i}+3\hat{j}+\lambda\hat{k}$ on $5\hat{i}-\hat{j}-3\hat{k}$, then λ is equal to

(1) ± 4 (2) ± 3 (3) ± 5 (4) ± 1

(22) If (1, 2, 4) and $(2, -3\lambda-3)$ are the initial and terminal points of the vector $\hat{i}+5\hat{j}-7\hat{k}$, then the value of λ is equal to

(1) $\frac{7}{3}$ (2) $-\frac{7}{3}$ (3) $-\frac{5}{3}$ (4) $\frac{5}{3}$

(23) If the points whose position vectors $10\hat{i}+3\hat{j}$, $12\hat{i}-5\hat{j}$ and $a\hat{i}+11\hat{j}$ are collinear then a is equal to

(1) 6 (2) 3 (3) 5 (4) 8

(24) If $\vec{a}=\hat{i}+\hat{j}+\hat{k}$, $\vec{b}=2\hat{i}+x\hat{j}+\hat{k}$, $\vec{c}=\hat{i}-\hat{j}+4\hat{k}$ and $\vec{a}\cdot(\vec{b}\times\vec{c})=70$, then x is equal to

(1) 5 (2) 7 (3) 26 (4) 10

(25) If $\vec{a}=\hat{i}+2\hat{j}+2\hat{k}, |\vec{b}|=5$ and the angle between $\vec{a}$ and $\vec{b}$ is $\frac{\pi}{6}$, then the area of the triangle formed by these two vectors as two sides, is

(1) $\frac{7}{4}$ (2) $\frac{15}{4}$ (3) $\frac{3}{4}$ (4) $\frac{17}{4}$

SUMMARY

In this chapter we have acquired the knowledge of the following :

- A **scalar** is a quantity that is determined by its magnitude.
- A **vector** is a quantity that is determined by both its magnitude and its direction
- If we have a liberty to choose the origins of the vector at any point then it is said to be a **free vector,** whereas if it is restricted to a certain specified point then the vector is said to be a **localized vector.**
- Two or more vectors are said to be **coplanar** if they lie on the same plane or parallel to the same plane.
- Two vectors are said to be **equal** if they have equal length and the same direction.
- A vector of magnitude 0 is called the **zero vector.**
- A vector of magnitude 1 is called a **unit vector**.

08_11th-Maths-Vol2-Unit8.indd 82 21/11/2024 15:02:49