www.tntextbooks.in

Evaluate Yourself

1) Draw the lewis structures for

   i) Nitrous acid (HNO_2)

   ii) Phosphoric acid

   iii) Sulphur troxide (SO_3)

10.2.3 Formal charge:

Let us draw the Lewis structure for carbon dioxide.

1. Skeletal structure

   O C O

2. Total number of valence electrons in CO_2

   = [1 x 4(carbon)] +[2 x 6(oxygen)] = 4+ 12 = 16

3. Draw single bonds between atoms. Two bonds can be drawn as shown in the figure for CO_2 which accounts for four electrons (2 bond pairs).

   O — C — O

4. Distribute the remaining twelve electrons (16 - 4= 12) as six lone pairs starting from most electronegative atom, the oxygen. Six lone pairs are distributed to the two terminal oxygens (three each) to satisfy their octet.

   :Ö — C — Ö:

5. Verify weather all the atoms have octet configuration. In the above distribution, the central carbon has two pair short for octet. Therefore, to satisfy the octet rule two lone pairs from one oxygen or one pair from each oxygen can be moved to form multiple bonds, leading the formation of two possible structures for carbon dioxide as shown below

   Ö = C = Ö :O ≡ C — Ö:

   structure 1 structure 2

Fig 10. 7 (a) two possible structures for carbon dioxide

Similarly, the Lewis structure for many molecules drawn using the above steps gives more than one acceptable structure. Let us consider the above mentioned two structures of carbon dioxide.

Which one the above forms represents the best distribution of electrons in the molecule. To find an answer, we need to know the formal charge of each atom in the Lewis structures. Formal charge of an atom in a molecule, is the electrical charge difference between the valence electron in an isolated atom and the number of electrons assigned to that atom in the Lewis structure.

$$\text{Formal charge of an atom} = N_v - \left(N_l + \frac{N_b}{2} \right)$$

Where,

N_v - Number of valence electron of atom in its isolated state.

N_l - Number of electrons present as lone pairs around the atom in the Lewis structure

N_b - Number of electrons present in bonds around the atom (bond pairs) in the