on. They are mostly chemoautotrophs. The unique feature of this group is the presence of lipids like glycerol & isopropyl ethers in their cell membrane. Due to the unique chemical composition the cell membrane show resistance against cell wall antibiotics and lytic agents. Example: *Methanobacterium, Halobacterium, Thermoplasma*.

1.4.9 Cyanobacteria (Blue Green Algae)

How old are Cyanobacteria ?
Stromatolites reveals the truth.



Stromatolites are deposits formed when colonies of cyanobacteria bind with calcium carbonate. They have a geological age of 2.7 billion years. Their abundance in the fossil record indicates that cyanobacteria helped in raising the level of free oxygen in the atmosphere.



- *Pseudomonas putida* is a superbug genetically engineered which breakdown hydrocarbons.
- "Pruteen" is a single cell protein derived from *Methylophilus methylotrophus*.
- *Agrobacterium tumefaciens* cause crown gall disease in plants but its inherent tumour inducing principle helps to carry the desired gene into the plant through Genetic engineering.
- *Thermus aquaticus* is a thermophilic gram negative bacteria which produces Taq Polymerase a key enzyme for Polymerase Chain Reaction (PCR).
- *Methanobacterium* is employed in biogas production. *Halobacterium*, an extremophilic bacterium grows in high salinity. It is exploited for the production β carotene.

Cyanobacteria are popularly called as 'Blue green algae' or 'Cyanophyceae'. They are photosynthetic, prokaryotic organisms. According to evolutionary record Cyanobacteria are primitive forms and are found in different habitats. Most of them are fresh water and few are marine (*Trichodesmium* and *Dermacarpa*) *Trichodesmium erythraeum* a cyanobacterium imparts red colour to Red sea. Species of *Nostoc*, *Anabaena* lead an endophytic life in the coralloid root of *Cycas*, leaves of aquatic fern *Azolla* by establishing a symbiotic association and fix atmospheric nitrogen. Members like *Gloeocapsa, Nostoc*, *Scytonema* are found as phycobionts in lichen thalli.

Salient features

- The members of this group are prokaryotes and lack motile reproductive structures.
- The thallus is unicellular in *Chroococcus*, Colonial in *Gloeocapsa* and filamentous trichome in *Nostoc*.
- Gliding movement is noticed in some species (*Oscillatoria*).
- The protoplasm is differentiated into central region called centroplasm and peripheral region bearing chromatophore called chromoplasm.
- The photosynthetic pigments include c-phyocyanin and c-phycoerythrin along with myxoxanthin and myxoxanthophyll.
- The reserve food material is Cyanophycean starch.
- In some forms a large colourless cell is found in the terminal or intercalary position called Heterocysts. They are involved in nitrogen fixation.
- They reproduce only through vegetative methods and produce Akinetes (thick wall dormant cell formed from vegetative cell), Hormogonia (a portion of filament get detached and reproduce by cell division), fission and endospores.
- The presence of mucilage around the thallus is characteristic feature of this group. Therefore, this group is also called Myxophyceae.
- Sexual reproduction is absent.
- *Microcystis aeruginosa*, *Anabaena flos-aquae* cause water blooms and release toxins and affect the aquatic organism.

11th Botany_EM_Chapter 1.indd 19 06/01/2023 19:44:55