



Amber is a plant secretion which is an efficient preservative that doesn't get degraded and hence can preserve remains of extinct life forms. The amber is produced by ***Pinites succinifera***, a Gymnosperm.

In this chapter we shall discuss in detail about one group of seed producing plants called **Gymnosperms**.

Gymnosperms (Gr. Gymnos = naked; sperma= seed) are naked seed producing plants. They were dominant in the Jurassic and Cretaceous periods of Mesozoic era. The members are distributed throughout the temperate and tropical region of the world

2.6.1 General characteristic features

- Most of the gymnosperms are evergreen, woody trees or shrubs. Some are lianas (*Gnetum*)
- The plant body is sporophyte and is differentiated into root, stem and leaves.
- A well developed tap root system is present. Coralloid roots of *Cycas* have symbiotic association with blue green algae. In *Pinus* the roots have mycorrhizae.
- The stem is aerial, erect and branched or unbranched (*Cycas*) with leaf scars.
- In conifers two types of branches namely branches of limited growth (Dwarf shoot) and Branches of unlimited growth (Long shoot) is present.
- Leaves are dimorphic, foliage and scale leaves are present. Foliage leaves are green, photosynthetic and borne on branches of limited growth. They show xerophytic features.
- The xylem consists of tracheids but in *Gnetum* and *Ephedra* vessels are present.
- Secondary growth is present. The wood may be **Manoxylic** (Porous, soft, more parenchyma with wide medullary ray -*Cycas*) or **Pycnoxylic** (compact with narrow medullary ray-*Pinus*).
- They are heterosporous. The plant may be monoecious (*Pinus*) or dioecious (*Cycas*).
- Microsporangia and megasporangia are produced on microsporophyll and megasporophyll respectively.
- Male and female cones are produced.
- Anemophilous pollination is present.
- Fertilization is siphonogamous and pollen tube helps in the transfer of male nuclei.
- Polyembryony (presence of many embryo) is present. The naked ovule develops into seed. The **endosperm** is haploid and develop before fertilization.
- The life cycle shows alternation of generation. The sporophytic phase is dominant and gametophytic phase is highly reduced. The photograph of some of the gymnosperms is given in Figure 2.22




a) *Taxus* b) *Ginkgo*

Figure 2.22: Gymnosperms

2.6.2 Classification of Gymnosperms

Sporne (1965) classified gymnosperms into 3 classes, 9 orders and 31 families. The classes include i) Cycadospsida ii) Coniferopsida iii) Gnetopsida.

GYMNOSPERMS		
Class-I	Class-II	Class-III
Cycadopsida	**Coniferopsida**	**Gnetopsida**
Orders: 1. Pteridosper-males 2. Bennettitales 3. Pentoxylales 4. Cycadales	Orders: 1. Cordaitales 2. Coniferales 3. Taxales 4. Ginkgoales	Order: 1. Gnetales

General Characters of Main classes:

Class I – Cycadopsida

- Plants are palm-like or fern-like.
- Compound, frond-like pinnate leaves.
- Manoxylic wood.

11th Botany_EM_Chapter 2.indd 51 06/01/2023 19:41:33