Palaeobotany in India

The National wood fossil park is situated in Tiruvakkarai, a Village of Villupuram district of Tamil Nadu. The park contains petrified wood fossils approximately 20 million years old. The term 'form genera' is used to name the fossil plants because the whole plant is not recovered as fossils instead organs or parts of the extinct plants are obtained in fragments. Shiwalik fossil park-Himachal Pradesh, Mandla Fossil park-Madhya Pradesh, Rajmahal Hills–Jharkhand, Ariyalur – Tamilnadu are some of the fossil rich sites of India.

Some of the fossil representatives of different plant groups are given below

Fossil Algae - *Palaeoporella*, *Dimorphosiphon*

Fossil Bryophytes – *Naiadita, Hepaticites, Muscites*

Fossil Pteridophytes – *Cooksonia, Rhynia,, Baragwanthia, Calamites*

Fossil Gymnosperms – *Medullosa, Lepido-carpon, Williamsonia, Lepidodendron*

Fossil Angiosperms – *Archaeanthus, Furcula*

Prof. Birbal Sahni (1891-1949)



Father of Indian Palaeobotany. He described Fossil plants from Rajmahal Hills of Eastern Bihar. *Pentoxylon sahnii*, *Nipanioxylon* are some of the form genera described by him. Birbal Sahni Institute of Palaebotany is located in Lucknow.

2.7 Angiosperms



In the previous section, the characteristic features of one of the spermatophyte called Gymnosperms were discussed. Spermatophytes also include plants bearing ovules enclosed in a protective cover called ovary, such plants are called Angiosperms. They constitute major plant group of our earth and are adapted to the terrestrial mode of life. This group of plants appeared during the early cretaceous period (140 million years ago) and dominates the vegetation on a global scale. The sporophyte is the dominant phase and gametophyte is highly reduced.

2.7.1 Salient features of Angiosperms

- Vascular tissue (Xylem and Phloem) is well developed.
- Flowers are produced instead of cone
- The Ovule remains enclosed in the ovary.
- Pollen tube helps in fertilization, so water is not essential for fertilization.
- Double fertilization is present. The endosperm is triploid.
- Angiosperms are broadly classified into two classes namely Dicotyledons and Monocotyledons.

2.7.2 Characteristic features of Dicotyledons and Monocotyledons

Dicotyledons

Morphological features

Reticulate venation is present in the leaves. Presence of two cotyledons in the seed. Primary root radicle persists as tap root.

Flowers tetramerous or pentamerous.

Tricolpate (3 furrow) pollen is present.

Anatomical features

- Vascular bundles are arranged in the form of a ring in stem.
- Vascular bundles are open (Cambium present).
- Secondary growth is present.

Monocotyledons

Morphological features

Parallel venation is present in the leaves. Presence of single cotyledon in the seed. Radicle doesn't persist and fibrous root is present.

11th Botany_EM_Chapter 2.indd 57 06/01/2023 19:41:40