

Figure 4.13: (c) Deciduous

c. Persistant: Calyx that persists and continues to be along with the fruit and forms a cup at the base of the fruit. Example: Brinjal.

d. Accrescent: Calyx that is persistent, grows along with the fruit and encloses the fruit either completely or partially. Example: *Physalis*.



Figure 4.13: (d) Persistant calyx



Figure 4.13: (e) Accrescent

3. Shapes of calyx

Have you noticed the shoe flower's calyx? It is bell shaped called **Campanulate**. The fruiting calyx is urn shaped in *Withania* and it is called **urceolate**. In *Datura* calyx is tube like and it is known as **tubular**. Two lipped calyx is present *in Ocimum*. Sometimes calyx is coloured and called **petaloid.** Example: *Saraca* and *Mussanda*. In Tridax, calyx is modified into hair like structures are called pappus.



Figure 4.14: (a) Companulate



Figure 4.14: (b) Pappus



Figure 4.14: (c) Mussaenda

4.3.3 Corolla

Corolla is the most attractive part in majority of the flowers and is usually brightly coloured. Corolla helps to display the flower and attracts the pollinators.

1. Fusion:

a. **Apopetalous (polypetalous)**: Petals are distinct. Example: *Hibiscus*.

b. **Sympetalous (gamopetalous)**: Petals are fused. Example: *Datura*.

4.3.4 Perianth

Can you recall the term homochlamydeous? Undifferentiated calyx and corolla in a flower is called **perianth**. Each member is called **tepal**. If the tepals are distinct they are called **Apotepalous** (Polyphyllous). Example: *Allium sativum*. Fused tepals are called **Syntepalous**. (Gamophyllous). Example: *Allium cepa*.

Lodicule: Reduced scale like perianth in the members of Poaceae is called lodicule.

4.3.5 Aestivation: Arrangement of sepals and petals in the flower bud is said to be aestivation.




Figure 4.15: Aestivation

11th Botany_EM_Chapter 4.indd 86 05-01-2023 18:22:58