of Taxonomy." He outlined an artificial system of classification in "***Species Plantarum***" in 1753, wherein he listed and described 7,300 species and arranged in 24 classes mostly on the basis of number, union (adhesion and cohesion), length, and distribution of stamens. The classes were further subdivided on the basis of carpel characteristics into orders. Hence the system of classification is also known as **sexual system of classification**.



Figure 5.4: Carolus Linnaeus

This system of classification though artificial, was continued for more than 100 years after the death of Linnaeus, due to its simplicity and easy way of identification of plants.

However the system could not hold good due to the following reasons.

1. Totally unrelated plants were kept in a single group, whereas closely related plants were placed in widely separated groups. Example:
   a. Zingiberaceae of monocotyledons and Anacardiaceae of dicotyledonous were placed under the class **Monandria** since these possess single stamens.
   b. *Prunus* was classified along with *Cactus* because of the same number of stamens.

No attempts were made to classify plants based on either natural or phylogenetic relationships which exist among plant groups.

5.10.2 Natural system

Botanists who came after Linnaeus realised that no single character is more important than the other characters. Accordingly an approach to a natural system of classification sprouted in France. The first scheme of classification based on overall similarities was presented by **Antoine Laurent de Jessieu** in 1789.

Bentham and Hooker system of classification



Figure 5.5: George Bentham and J.D. Hooker

A widely followed natural system of classification considered the best was proposed by two English botanist **George Bentham** (1800 – 1884) and **Joseph Dalton Hooker** (1817–1911). The classification was published in a three volume work as "*Genera Plantarum*" (1862–1883) describing 202 families and 7569 genera and 97,205 species. In this system the seeded plants were classified into 3 major classes such as Dicotyledonae, Gymnospermae and Monocotyledonae.

Class I Dicotyledonae: Plants contain two cotyledons in their seed, leaves with reticulate venation, tap root system and tetramerous or pentamerous flowers come under this class. It includes three sub-classes – **Polypetalae, Gamopetalae** and **Monochlamydeae**.

Sub-class 1. Polypetalae: Plants with free petals and dichlamydeous flowers come under polypetalae. It is further divided into three series – **Thalamiflorae, Disciflorae** and **Calyciflorae**.

Series (i) Thalamiflorae: Plants having flowers with dome or conical shaped thalamus and superior ovary are included in this series. It includes 6 orders and 34 families.

Series (ii) Disciflorae: Flowers having prominent disc shaped thalamus with superior ovary come under this series. It includes 4 orders and 23 families.

Series (iii) Calyciflorae: It includes plants having flowers with cup shaped thalamus and

11th Botany_EM_Chapter 5.indd 110 06/01/2023 19:43:37