


Figure 5.10: Diagramatic representation of class Magnoliopsida and Liliopsida.

5.10.4 Angiosperm phylogeny group (APG) classification

The most recent classification of flowering plants based on **phylogenetic data** was set in the last decade of twentieth century. Four versions of Angiosperm Phylogenetic Group classification (APG I, APG II, APG III & APG IV) have been published in 1998, 2003, 2009 and 2016 respectively. Each version supplants the previous version. Recognition of **monophyletic** group based on the information received from various disciplines such as gross morphology, anatomy, embryology, palynology, karyology, phytochemistry and more strongly on molecular data with respect to DNA sequences of two chloroplast genes (*atp*B and *rbc*L) and one nuclear gene (nuclear ribosomal 18s DNA).

The most recent updated version, APG IV (2016) recognised 64 orders and 416 families. Of these, 416 families 259 are represented in India.

The outline of APG IV classification is given below.

Angiosperms are classified into three clades early angiosperms, monocots and eudicots. Early angiosperms are classified into **8 orders** and **26 families** (**ANA**-clade + magnoliids + Chloranthales)




Figure 5.11: Simplified version of APG IV
(Source: Plant Gateway's The Global Flora, Vol. I January 2018)

Amborellales
Nymphaeales
Austrobaileyales

- Seeds generally always with two cotyledons.

11th Botany_EM_Chapter 5.indd 113 06/01/2023 19:43:39