plants." Angiosperm phylogeny group (APG) classification is the most recent classification of flowering plants based on phylogenetic data. APG system is an evolving and currently accepted system across the world and followed by all the leading taxonomic institutions and practising taxonomists.

Cladistics is the methodology, used to classify organisms into monophyletic groups, consisting of all the descents of the common ancestors. The outcome of a cladistic analysis is a cladogram and is constructed to represent the best hypothesis of phylogenetic relationships. Chemotaxonomy is the scientific approach of classification of plants on the basis of their biochemical constituents in them. Utilization of the characters of chromosome for the taxonomic inference is known as karyotaxonomy. The application of serology in solving taxonomic problems is called serotaxonomy. Molecular Taxonomy is the branch of phylogeny that analyses hereditary molecular differences, mainly in DNA nuclear and chloroplast sequences, to gain information and to establish genetic relationship between the members of different taxonomic categories. Different molecular markers like allozymes, mitochondrial DNA, microsatellites, RAPDs, AFLPs, single nucleotide polymorphism-(SNP), microchips or arrays are used in analysis. Molecular Taxonomy unlocks the treasure chest of information on evolutionary history of organisms. It plays a vital role in phytogeography, which ultimately helps in genome mapping and biodiversity conservation. DNA barcoding is a taxonomic method that uses a very short genetic sequence from a standard part of a genome. It helps in identification of organisms.

Evaluation

HXN9Q

1. Phylogenetic classification is the most favoured classification because it reflects
   a. Comparative Anatomy
   b. Number of flowers produced
   c. Comparative cytology
   d. Evolutionary relationships
2. The taxonomy which involves the similarities and dissimilarities among the immune system of different taxa is termed as
   a. Chemotaxonomy
   b. Molecular systematics
   c. Serotaxonomy
   d. Numerical taxonomy
3. Perianth is present in
   a. *Clitoria ternatea*
   b. *Datura metal*
   c. *Allium cepa*
   d. *Pongamia pinnata*
4. Flowers are zygomorphic in
   a. *Ceropegia* b. *Thevetia*
   c. *Datura* d. *Solanum*
5. What is the role of national gardens in conserving biodiversity – discuss
6. Where will you place the plants which contain two cotyledons with cup shaped thalamus?
7. Write any five economic importance of family Euphorbiaceae.
8. How will you distinguish Solanaceae members from Liliaceae members?
9. Write the botanical description of *Clitoria ternatea.*
10. Write the botanical description of *Musa paradisiaca.*

11th Botany_EM_Chapter 5.indd 142 06/01/2023 19:43:57