


Figure 6.16: Structure of Chloroplast




Figure 6.17: Structure of Ribosomes



Svedberg unit (s).

The size of ribosomes and their subunits are usually given in Svedberg unit (named after Theoder Svedberg, Swedish Chemist Noble Laureate 1929), a measure of a particle size dependent on the speed with which particle sediment in the ultracentrifuge.

Ribosome consists of RNA and protein: RNA 60 % and protein 40%. During protein synthesis, many ribosomes are attached to the single mRNA and is called **polysomes** or **polyribosomes**. The function of polysomes is the formation of several copies of a particular polypeptide during protein synthesis. They are free in non-protein synthesising cells. In protein synthesising cells they are linked together with the help of Mg^{2+} ions.

Types of Ribosomes

70S Ribosomes (sub unit 30S and 50S)	80S Ribosomes (sub units 40S and 60S)
3 RNA molecule (i) 16SrRNA in 30S subunit (ii) 23S and 5S in 50S large subunit **(Prokaryotic cells of bluegreen algae bacteria, mitochondria and chloroplast of many algae and higher plants)**	4 RNA molecule (i) 18SrRNA in 40S small subunit (ii) 28S, 5.8S and 5S in larger 60S subunit **(Eukaryotic cells of plants and animals)**

6.6.8 Lysosomes (Suicidal Bags of Cell)

Lysosomes were discovered by **Christian de Duve** (1953), these are known as **suicidal bags**. They are spherical bodies enclosed by a single unit membrane. They are found in eukaryotic cell. Lysosomes are small vacuoles formed when small pieces of golgi body are pinched off from its tubules.

11th Botany_EM_Chapter 6.indd 157 05-01-2023 18:25:13