(Figure 11.18). This mass of tissue lies near vein endings (xylem and Phloem). The liquid coming out of hydathode is not pure water but a solution containing a number of dissolved substances.

11.6.7 Measurement of Transpiration

1. Ganongs potometer

Ganongs potometer is used to measure the rate of transpiration indirectly. In this, the amount of water absorbed is measured and assumed that this amount is equal to the amount of water transpired.

Apparatus consists of a horizontal graduated tube which is bent in opposite directions at the ends. One bent end is wide and the other is narrow. A reservoir is fixed to the horizontal tube near the wider end. The reservoir has a stopcock to regulate water flow. The apparatus is filled with water from reservoir. A twig or a small plant is fixed to the wider arm through a split cock. The other bent end of the horizontal tube is dipped into a beaker containing coloured water. An air bubble is introduced into the graduated tube at the narrow end (Figure 11.19). keep this apparatus in bright sunlight and observe.As transpiration takes place, the air bubble will move towards the twig. The loss is compensated by water absorption through the xylem portion of the twig. Thus, the rate of water absorption is equal to the rate of transpiration.




Figure 11.19: Ganongs Potometer

2. Cobalt chloride ($CoCl_2$) paper method

Select a healthy dorsiventral leaf and clean its upper and lower surface with dry cotton. Now place a dry Cobalt chloride (**$CoCl_2$**) strips on both surface and immediately cover the paper with glass slides and immobilize them. It will be observed after some time that the $CoCl_2$ strip of lower epidermis turns pink. This indicates that $CoCl_2$ becomes hydrated ($CoCl_2 \cdot 2H_2O$ or $CoCl_2 \cdot 4H_2O$) due to water vapours coming out through stomata. The rate of transpiration is more on the lower surface than in the upper surface of the dorsiventral leaf.

11.6.8 Significance of transpiration

Transpiration leads to loss of water, as stated earlier in this lesson 95% of absorbed water is lost in transpiration. It seems to be an evil process to plants. However, number of process like absorption of water, ascent of sap and mineral absorption directly rely on the transpiration. Moreover plants withstand against scorching sunlight due to transpiration. Hence the transpiration is a "**necessary evil**" as stated by **Curtis**.

11.7 Translocation of Organic Solutes

Leaves synthesize food material through photosynthesis and store in the form of starch grains. When required the starch is converted into simple sugars. They must be transported to various parts of the plant system for further utilization. However, the site of food production (leaves) and site of utilization are separated far apart. Hence, the organic food has to be transported to these areas.

The phenomenon of food transportation from the site of synthesis to the site of utilization is known as **translocation of organic solutes**. The term **solute** denotes food material that moves in a solution.

11.7.1 Path of Translocation

It has now been well established that phloem is the path of translocation of solutes. Ringing or girdling experiment will clearly demonstrate the translocation of solute by phloem.

XIth_Botany_Chapter_11_EM.indd 251 05-01-2023 18:29:10