gibberellin by **Yabuta** (1935). These are more than 100 gibberellins reported from both fungi and higher plants. They are noted as GA_1, GA_2, GA_3 and so on. GA_3 is the first discovered gibberellin. In 1938, **Yabuta** and **Sumiki** isolated gibberellin in crystalline form. In1955, **Brain** *et al.,* gave the name **gibberellic acid**. In 1961, Cross *et al.,* established its structure.

Agent Orange
Mixture of two phenoxy herbicides 2,4-D and 2,4,5-T is given the name 'Agent orange' which was used by USA in Vietnam war for defoliation of forest (chemical warfare).



In botanical gardens and tea gardens, gardeners trim the plants regularly so that they remain bushy. Does this practice have any scientific explanation?

Yes, trimming of plants removes apical buds and hence apical dominance. The lateral buds sprout and make the plants bushy.

2. Occurrence
The major site of gibberellin production in plants is parts like embryo, roots and young leaves near the tip. Immature seeds are rich in gibberellins.

3. Precursors
The gibberellins are chemically related to terpenoids (natural rubber, carotenoids and steroids) formed by 5-C precursor, an Isoprenoid unit called Iso Pentenyl Pyrophosphate (IPP) through a number of intermediates. The primary precursor is acetate.

4. Chemical structure
All gibberellins have gibbane ring structure.

5. Transport in plants
The transport of gibberellins in plants is non-polar. Gibberellins are translocated through phloem and also occur in xylem due to lateral movement between vascular bundles.

6. Bioassay (Dwarf Pea assay)
Seeds of dwarf pea are allowed to germinate till the formation of the coleoptile. GA solution is applied to some seedlings. Others are kept under control. Epicotyl length is measured and as such, GA stimulating epicotyl growth can be seen.

7. Physiological Effects
- It produces extraordinary elongation of stem caused by cell division and cell elongation.
- Rosette plants (genetic dwarfism) exhibit excessive internodal growth when they are treated with gibberellins. This sudden elongation of stem followed by flowering by the application of gibberellin is called bolting (Figure 15.9).
- Gibberellin breaks dormancy in potato tubers.
- Many biennials usually flower during second year of their growth. For flowering in the first year it self these plants should be treated with gibberellins.
- Formation of seedless fruits without fertilization is induced by gibberellins Example: Seedless tomato, apple and cucumber.
- Promotes elongation of inter-node in sugarcane without decreasing sugar content.
- Promotion of flowering in long day plants even under short day conditions.
- It stimulates the seed germination.




Figure 15.9: Bolting

15.2.3 Cytokinins (Cytos – cell, Kinesis – division)

1. Discovery
The presence of cell division inducing substances in plants was first demonstrated by **Haberlandt** in 1913 in Coconut milk (liquid

11th Botany_EM_Chapter 15.indd 312 11/01/2023 15:00:25