where n is the number of moles of gas and V is the volume of the container

$$\Rightarrow P' \alpha \frac{n^2}{V^2}$$

$$\Rightarrow P' = a \frac{n^2}{V^2}$$

where a is proportionality constant and depends on the nature of gas

Therefore,

$$P_{ideal} = P + \frac{an^2}{V^2} \quad \text{------ (6.20)}$$

Volume Correction

As every individual molecule of a gas occupies a certain volume, the actual volume is less than the volume of the container, V. Van der Waals introduced a correction factor V' to this effect. Let us calculate the correction term by considering gas molecules as spheres.




Figure. 6.11 Excluded volume

V = excluded volume

Excluded volume for two molecules

$$= \frac{4}{3}\pi(2r)^3$$

$$= 8\left(\frac{4}{3}\pi r^3\right) = 8\ V_m$$

where v_m is a volume of a single molecule

Excluded volume for single molecule

$$= \frac{8\ V_m}{2} = 4\ V_m$$

Excluded volume for n molecule

$$= n\ (4\ V_m) = nb$$

Where b is van der waals constant whch is equal to $4\ V_m$

$$\Rightarrow V' = nb$$

$$V_{ideal} = V - nb \text{ ----- (6.21)}$$

Replacing the corrected pressure and volume in the ideal gas equation PV=nRT we get the van der Waals equation of state for real gases as below,

$$\left(P + \frac{an^2}{V^2}\right)(V - nb) = nRT \text{ . ----(6.22)}$$

The constants *a* and *b* are van der Waals constants and their values vary with the nature of the gas. It is an approximate formula for the non-ideal gas.

6.6 Pressure-Volume isotherms of Carbon dioxide

Andrew's isotherm

Thomas Andrew gave the first complete data on pressure-volume-temperature of a substance in the gaseous and liquid states. He plotted isotherms of carbon dioxide at different temperatures which is shown in Figure. 6.12. From the plots we can infer the following.

11th Std Chemistry 158-185.indd 172 23-12-2021 15:23:07