1.8.2 Dimensional Quantities, Dimensionless Quantities, Principle of Homogeneity

On the basis of dimension, we can classify quantities into four categories.

(1) **Dimensional variables**

Physical quantities, which possess dimensions and have variable values are called dimensional variables. Examples are length, velocity, and acceleration etc.

(2) **Dimensionless variables**

Physical quantities which have no dimensions, but have variable values are called dimensionless variables. Examples are specific gravity, strain, refractive index etc.

(3) **Dimensional Constant**

Physical quantities which possess dimensions and have constant values are called dimensional constants. Examples are Gravitational constant, Planck's constant etc.

(4) **Dimensionless Constant**

Quantities which have constant values and also have no dimensions are called dimensionless constants. Examples are π, e (Euler's number), numbers etc.

Principle of homogeneity of dimensions

The principle of homogeneity of dimensions states that the dimensions of all the terms in a physical expression should be the same. For example, in the physical expression $v^2 = u^2 + 2as$, the dimensions of v^2, u^2 and 2 as are the same and equal to $[L^2T^{-2}]$.

1.8.3 Application and Limitations of the Method of Dimensional Analysis.

This method is used to

(i) Convert a physical quantity from one system of units to another.

(ii) Check the dimensional correctness of a given physical equation.

(iii) Establish relations among various physical quantities.

(i) To convert a physical quantity from one system of units to another

This is based on the fact that the product of the numerical values (n) and its corresponding unit (u) is a constant. i.e, n [u] = constant (or) $n_1[u_1] = n_2[u_2]$.

Consider a physical quantity which has dimension '*a*' in mass, '*b*' in length and '*c*' in time. If the fundamental units in one system are M_1, L_1 and T_1 and the other system are M_2, L_2 and T_2 respectively, then we can write, $n_1 [M_1^a L_1^b T_1^c] = n_2 [M_2^a L_2^b T_2^c]$

We have thus converted the numerical value of physical quantity from one system of units into the other system.

EXAMPLE 1.12

Convert 76 cm of mercury pressure into Nm^{-2} using the method of dimensions.

Solution

In cgs system 76 cm of mercury pressure = $76 \times 13.6 \times 980$ dyne cm^{-2}

The dimensional formula of pressure P is $[ML^{-1}T^{-2}]$

UNIT-1(XI-Physics_Vol-1).indd 32 05-01-2022 22:12:00