EXAMPLE 2.37

Suppose an object is thrown with initial speed 10 m s^{-1} at an angle $\pi/4$ with the horizontal, what is the range covered? Suppose the same object is thrown similarly in the Moon, will there be any change in the range? If yes, what is the change? (The acceleration due to gravity in the Moon $g_{moon} = \frac{1}{6} g$)

Solution

In projectile motion, the range of particle is given by,

$$R = \frac{u^2 \sin 2\theta}{g}$$

$$\theta = \pi/4 \quad u = v_0 = 10 m\ s^{-1}$$

$$\therefore R_{earth} = \frac{(10)^2 \sin \pi/2}{9.8} = 100/9.8$$

$$R_{earth} = 10.20\ m \text{ (Approximately 10 m)}$$

If the same object is thrown in the Moon, the range will increase because in the Moon, the acceleration due to gravity is smaller than g on Earth,

$$g_{moon} = \frac{g}{6}$$

$$R_{moon} = \frac{u^2 \sin 2\theta}{g_{moon}} = \frac{v_0^{\,2} \sin 2\theta}{g/6}$$

$$\therefore R_{moon} = 6R_{earth}$$


$$R_{moon} = 6 \times 10.20 = 61.20\ m$$


(Approximately 60 m)

The range attained on the Moon is approximately six times that on Earth.

EXAMPLE 2.38

In the cricket game, a batsman strikes the ball such that it moves with the speed 30 m s^{-1} at an angle 30^0 with the horizontal as shown in the figure. The boundary line of the cricket ground is located at a distance of 75 m from the batsman? Will the ball go for a six? (Neglect the air resistance and take acceleration due to gravity g = 10 m s^{-2}).





Solution

The motion of the cricket ball in air is essentially a projectile motion. As we have already seen, the range (horizontal distance) of the projectile motion is given by

$$R = \frac{u^2 \sin 2\theta}{g}$$

The initial speed $u = 30\ m\ s^{-1}$

The projection angle $\theta = 30°$

UNIT-2(XI-Physics_Vol-1)-1.indd 87 05-01-2022 22:15:43