**Tangential acceleration**

Consider an object moving along a circle of radius r. In a time $\Delta t$, the object travels an arc distance $\Delta s$ as shown in Figure 2.47. The corresponding angle subtended is $\Delta\theta$





**Figure 2.47** Circular motion

The $\Delta s$ can be written in terms of $\Delta\theta$ as,

$$\Delta s = r\Delta\theta \qquad (2.35)$$

In a time $\Delta t$, we have

$$\frac{\Delta s}{\Delta t} = r\frac{\Delta\theta}{\Delta t} \qquad (2.36)$$

In the limit $\Delta t \to 0$, the above equation becomes

$$\frac{ds}{dt} = r\omega \qquad (2.37)$$

Here $\frac{ds}{dt}$ is linear speed $(v)$ which is tangential to the circle and $\omega$ is angular speed. So equation (2.37) becomes

$$v = r\omega \qquad (2.38)$$

which gives the relation between linear speed and angular speed.

**Note**

**The direction of linear velocity $\vec{v}$ is tangential to the circle, whereas the direction of angular velocity $\vec{\omega}$ is along the axis of rotation. The radius is also represented as a vector $\vec{r}$ directed radially from the centre of the circle.**

Equation (2.38) is true only for circular motion. In general the relation between linear and angular velocity is given by

$$\vec{v} = \vec{\omega} \times \vec{r} \qquad (2.39)$$

For circular motion equation (2.39) reduces to equation (2.38) since $\vec{\omega}$ and $\vec{r}$ are perpendicular to each other.

Differentiating the equation (2.38) with respect to time, we get (since r is constant)

$$\frac{dv}{dt} = \frac{rd\omega}{dt} = r\alpha \qquad (2.40)$$

Here $\frac{dv}{dt}$ is the tangential acceleration and is denoted as $a_t$. $\frac{d\omega}{dt}$ is the angular acceleration $\alpha$. Then eqn. (2.39) becomes

$$a_t = r\alpha \qquad (2.41)$$

The tangential acceleration $a_t$ experienced by an object in circular motion is shown in Figure 2.48.



UNIT-2(XI-Physics_Vol-1)-1.indd 90 05-01-2022 22:15:49