But

$$\int_A^C \vec{F}.d\vec{r} = \int_A^C \left(-mg\hat{j}\right).\left(dx\hat{i}\right) = 0$$

$$\int_C^D \vec{F}.d\vec{r} = \int_C^D \left(-mg\hat{j}\right).\left(dy\hat{j}\right)$$

$$= -mg\int_0^h dy = -mgh$$

$$\int_D^B \vec{F}.d\vec{r} = \int_D^B \left(-mg\hat{j}\right).\left(-dx\hat{i}\right) = 0$$

Therefore, the total work done by the force along the path 2 is

$$W_{\text{path 2}} = \int_A^B \vec{F}.d\vec{r} = -mgh$$

Note that the work done by the conservative force is independent of the path.

EXAMPLE 4.12

Consider an object of mass 2 kg moved by an external force 20 N in a surface having coefficient of kinetic friction 0.9 to a distance 10 m. What is the work done by the external force and kinetic friction ? Comment on the result. (Assume g = 10 ms^{-2})

Solution

m = 2 kg, d = 10 m, F_{ext} = 20 N, μ_k = 0.9.
When an object is in motion on the horizontal surface, it experiences two forces.

(a) External force, $F_{ext} = 20$ N

(b) Kinetic friction,
$f_k = \mu_k mg = 0.9\times(2)\times 10 = 18\,\text{N}$.

The work done by the external force $W_{ext} = F\,d = 20\times 10 = 200\,\text{J}$

The work done by the force of kinetic friction $W_k = f_k d = (-18)\times 10 = -180$ J. Here the negative sign implies that the force of kinetic friction is opposite to the direction of displacement.

The total work done on the object $W_{total} = W_{ext} + W_k = 200\text{ J} - 180\text{ J} = 20\text{ J}$.

Since the friction is a non-conservative force, out of 200 J given by the external force, the 180 J is lost and it can not be recovered.

4.2.8 Law of conservation of energy

When an object is thrown upwards its kinetic energy goes on decreasing and consequently its potential energy keeps increasing (neglecting air resistance). When it reaches the highest point its energy is completely potential. Similarly, when the object falls back from a height its kinetic energy increases whereas its potential energy decreases. When it touches the ground its energy is completely kinetic. At the intermediate points the energy is both kinetic and potential as shown in Figure 4.13. When the body reaches the ground the kinetic energy is completely dissipated into some other form of energy like sound, heat, light and deformation of the body etc.

In this example the energy transformation takes place at every point. The sum of kinetic energy and potential energy i.e., the total mechanical energy always remains constant, implying that the total energy is conserved. This is stated as the law of conservation of energy.

UNIT-4(XI-Physics_Vol-1).indd 183 05-01-2022 22:19:55