

Table 5.3 Moment of Inertia of Different Rigid Bodies

No.	Object	About an axis	Diagram	Moment of Inertia (I) kg m²	Radius of Gyration (K)	Ratio $\left(\frac{K^2}{R^2}\right)$
1.	**Thin Uniform Rod** Mass = M Length = ℓ	Passing through the centre and perpendicular to the length		$\frac{1}{12}M\ell^2$	$\frac{\ell}{\sqrt{12}}$	--
		Touching one end and perpendicular to the length		$\frac{1}{3}M\ell^2$	$\frac{\ell}{\sqrt{3}}$	
2.	**Thin Uniform Rectangular Sheet** Mass = M; Length = ℓ; Breadth = b	Passing through the centre and perpendicular to the plane of the sheet		$\frac{1}{12}M(\ell^2+b^2)$	$\sqrt{\frac{(\ell^2+b^2)}{12}}$	--
3.	Thin Uniform Ring Mass = M Radius = R	Passing through the centre and perpendicular to the plane		MR^2	R	1
		Touching the edge perpendicular to the plane (perpendicular tangent)		$2MR^2$	$\left(\sqrt{2}\right)R$	2
		Passing through the centre lying on the plane (along diameter)		$\frac{1}{2}MR^2$	$\left(\frac{1}{\sqrt{2}}\right)R$	$\frac{1}{2}$
		Touching the edge parallel to the plane (parallel tangent)		$\frac{3}{2}MR^2$	$\left(\sqrt{\frac{3}{2}}\right)R$	$\frac{3}{2}$
4.	**Thin Uniform Disc** Mass = M Radius = R	Passing through the centre and perpendicular to the plane		$\frac{1}{2}MR^2$	$\left(\frac{1}{\sqrt{2}}\right)R$	$\frac{1}{2}$
		Touching the edge perpendicular to the plane (perpendicular tangent to the plane)		$\frac{3}{2}MR^2$	$\left(\sqrt{\frac{3}{2}}\right)R$	$\frac{3}{2}$
		Passing through the centre lying on the plane (along diameter)		$\frac{1}{4}MR^2$	$\left(\frac{1}{2}\right)R$	$\frac{1}{4}$
		Touching the edge parallel to the plane (parallel tangent to the plane)		$\frac{5}{4}MR^2$	$\left(\sqrt{\frac{5}{4}}\right)R$	$\frac{5}{4}$

UNIT-5(XI-Physics_Vol-1).indd 244 06-01-2022 17:47:41