
When $m_1 = m_2 = m$

$$\vec{E}_{total} = -\frac{Gm}{a^2}(\hat{i} + \hat{j})$$

$(\hat{i} + \hat{j} = \hat{j} + \hat{i}$ as vectors obeys commutation law).




$\vec{E}_{total}$ points towards the origin of the co-ordinate system and the magnitude of $\vec{E}_{total}$ is $\sqrt{2}\frac{Gm}{a^2}$.

EXAMPLE 6.4

Qualitatively indicate the gravitational field of Sun on Mercury, Earth, and Jupiter shown in figure.

Since the gravitational field decreases as distance increases, Jupiter experiences a weak gravitational field due to the Sun. Since Mercury is the nearest to the Sun, it experiences the strongest gravitational field.



Solar System

6.2.3 Gravitational Potential Energy

The concept of potential energy and its physical meaning were dealt in unit 4. The gravitational force is a conservative force and hence we can define a gravitational potential energy associated with this conservative force field.

Two masses m_1 and m_2 are initially separated by a distance r'. Assuming m_1 to be fixed in its position, work must be done on m_2 to move the distance from r' to r as shown in Figure 6.12(a)




Figure 6.12 Two distant masses changing the linear distance

To move the mass m_2 through an infinitesimal displacement $d\vec{r}$ from $\vec{r}$ to $\vec{r} + d\vec{r}$ (shown in the Figure 6.12(b)), work has to be done externally. This infinitesimal work is given by

$$dW = \vec{F}_{ext} . d\vec{r} \qquad (6.21)$$

The work is done against the gravitational force, therefore,

$$\left|\vec{F}_{ext}\right| = \left|\vec{F}_G\right| = \frac{Gm_1m_2}{r^2} \qquad (6.22)$$

Substituting Equation (6.22) in 6.21, we get

$$dW = \frac{Gm_1m_2}{r^2}\hat{r} . d\vec{r} \qquad (6.23)$$

UNIT-6(XI-Physics_Vol-2).indd 13 02-01-2023 14:28:50