


Figure 7.17 (a) turbulent flow around a sphere (when $v = v_c$) (b) turbulent flow around a sphere (when $v > v_c$)

The distinction between the two types of motion can be easily demonstrated by injecting a jet of ink axially in a wide tube through which water flows. When the velocity of the fluid is small, the ink will move in a straight line path. Conversely, when the velocity is increased beyond a certain value, the ink will spread out showing the disorderliness and hence the motion becomes turbulent. The zig-zag motion results in the formation of eddy currents and as a consequence, much energy is dissipated.

7.4.4 Reynold's number

We have learnt that the flow of a fluid becomes steady or laminar when the velocity of flow is less than the critical velocity v_c otherwise, the flow becomes turbulent. Osborne Reynolds (1842-1912) formulated an equation to find out the nature of the flow of fluid, whether it is streamlined or turbulent.

$$R_c = \frac{\rho v D}{\eta} \qquad (7.21)$$

It is a dimensionless number called '*Reynold's number*'. It is denoted by the symbol R_c or K. In the equation, ρ denotes the density of the fluid, v the velocity of the flowing fluid, D is the diameter of the pipe in which the fluid flow, and η is the coefficient of viscosity of the fluid. The value of R_c remains the same in any system of units.

Table 7.3 To understand the flow of liquid, Reynold has estimated the value of R_c as follows

S. No.	Reynold's number	Flow
1	$R_c < 1000$	Streamline
2	$1000 < R_c < 2000$	Unsteady
3	$R_c > 2000$	Turbulent

Hence, Reynold's number R_c is a critical variable which decides whether the flow of a fluid through a cylindrical pipe is streamlined or turbulent.

In fact, the critical value of R_c at which the turbulence sets is found to be the same for geometrically similar flows. For example, when two liquids (say oil and water) of different densities and viscosities flow in pipes of same shapes and sizes, the turbulence sets in at almost the same value of R_c. The above fact leads to the *Law of similarity* which states that when there are two geometrically similar flows, both are essentially equal to each other, as long as they embrace the same Reynold's number. The *Law of similarity* plays a very important role in technological applications. The

UNIT-7(XI-Physics_Vol-2).indd 68 02-01-2023 14:39:56