not a quantity and through the work energy is transferred to the system . So we cannot use the word 'the object contains more work' or 'less work'.

Either the system can transfer energy to the surrounding by doing work on surrounding or the surrounding may transfer energy to the system by doing work on the system. For the transfer of energy from one body to another body through the process of work, they need not be at different temperatures.

8.1.4 Meaning of temperature

Temperature is the degree of hotness or coolness of a body. Hotter the body higher is its temperature. The temperature will determine the direction of heat flow when two bodies are in thermal contact.

The SI unit of temperature is ***kelvin*** **(K)**.

In our day to day applications, ***Celsius*** (˚C) and ***Fahrenheit*** (°F) scales are used.

Temperature is measured with a thermometer.

The conversion of temperature from one scale to other scale is given in Table 8.1

Table 8.1 Temperature conversion

Scale	To Kelvin	From Kelvin
Celsius	K=°C + 273.15	°C = K – 273.15
Fahrenheit	K=(°F + 459.67)÷1.8	°F=(K × 1.8)-459.67

Scale	To Fahrenheit	From Fahrenheit
Celsius	°F=(1.8 × °C) + 32	°C = (°F – 32)÷1.8

Scale	To Celsius	From Celsius
Fahrenheit	°C=(°F – 32)÷1.8	°F = (1.8 × °C) + 32

8.2 THERMAL PROPERTIES OF MATTER

8.2.1 Boyle's law, Charles' law and ideal gas law

For a given gas at low pressure (density) kept in a container of volume V, experiments revealed the following information.

- When the gas is kept at constant temperature, the pressure of the gas is inversely proportional to the volume. $P \propto \frac{1}{V}$. It was discovered by Robert Boyle (1627-1691) and is known as Boyle's law.
- When the gas is kept at constant pressure, the volume of the gas is directly proportional to absolute temperature. $V \propto T$. It was discovered by Jacques Charles (1743-1823) and is known as Charles' law.
- By combining these two equations we have

$PV = CT$. Here C is a positive constant.

We can infer that C is proportional to the number of particles in the gas container by considering the following argument. If we take two containers of same type of gas with same volume V, same pressure P and same temperature T, then the gas in each container obeys the above equation. $PV = CT$. If the two containers of gas is considered as a single system, then the pressure and temperature of this combined system will be same but volume will be twice and number of particles will also be double as shown in Figure 8.2

UNIT-8(XI-Physics_Vol-2).indd 97 02-01-2023 15:25:14