Trial No.	Microscope reading for the position of Lower meniscus of liquid			Microscope reading for the position of Lower tip of the needle			Height of the liquid h $\times 10^{-2}$ m a ~ b
	MSR	VSC × LC = VSR	TR (a) $\times 10^{-2}$ m	MSR	VSC × LC = VSR	TR (b) $\times 10^{-2}$ m	
						Mean h =	

Radius of the capillary tube

Tube	Microscope reading for the position of inner left wall of the tube L_1			Microscope reading for the position of inner right wall of the tube R_1			Radius of the capillary tube $r = \frac{1}{2}(L_1 \sim R_1)$ $\times 10^{-2}$ m
	MSR	VSC × LC = VSR	TR $\times 10^{-2}$ m	MSR	VSC × LC = VSR	TR $\times 10^{-2}$ m	

CALCULATION

Mean rise of the liquid in the capillary tube $h =$ _____ $\times 10^{-2}$ m

Diameter of the capillary tube $2r =$ _____ $\times 10^{-2}$ m

Radius of the capillary tube $r =$ _____ $\times 10^{-2}$ m

Density of the liquid $\sigma = 1000\ kg\ m^{-3}$

Acceleration due to gravity $g = 9.8\ m\ s^{-2}$

Surface tension $T = \frac{hr\sigma g}{2}$

$=$_____ $N\ m^{-1}$

RESULT

Surface tension of the given liquid by capillary rise method $T =$_____ $N\ m^{-1}$

UNIT-12(XI-Physics_Vol-2) PRACTICAL FIRST YEAR.indd 299 02-01-2023 15:28:05