into squamous epithelium, cuboidal epithelium, columnar epithelium, ciliated epithelium and pseudostratified epithelium (Figure 3.2). The **squamous epithelium** is made of a single thin layer of flattened cells with irregular boundaries. They are found in the kidney glomeruli, air sacs of lungs, lining of heart, blood vessels and lymphatic vessels and are involved in functions like forming a diffusion boundary and filtration in sites where protection is not important. The **cuboidal epithelium** is made of a single layer of cube like cells. This tissue is commonly found in the kidney tubules, ducts and secretory portions of small glands and surface of the ovary. Its main functions are secretion and absorption. The **columnar epithelium** is composed of single layer of tall cells with round to oval nuclei at the base. It lines the digestive tract from the stomach to the rectum. The two modifications of this lining are the presence of **microvilli** on the apical surface of the absorptive cells and **Goblet cell** which secretes the protective lubricating mucus. The functions of this epithelium include absorption, secretion of mucus, enzymes and other substances. If the columnar cells bear cilia on their free surfaces they are called **ciliated epithelium**. This **ciliated type** propels mucus by ciliary actions and it lines the small bronchioles, fallopian tubes and uterus. **Nonciliated type** lines most of the digestive tract, gall bladder and secretory ducts of glands.

Pseudo-stratified epithelial cells are columnar, but unequal in size. Although the epithelium is single layered yet it appears to be multi-layered because the nuclei lie at different levels in different cells. Hence, it is also called pseudostratified epithelium and




Figure 3.2 Types of Epithelial tissues

XI Std Zoology Chapter-3 EM.indd 39 30/12/2022 10:33:23