Nutrients like amino acids, glucose and electrolytes like Na^+ are absorbed into the blood against the concentration gradient by active transport. The insoluble substances like fatty acids, glycerol and fat soluble vitamins are first incorporated into small, spherical water soluble droplets called micelles and are absorbed into the intestinal mucosa where they are re-synthesized into protein coated fat globules called chylomicrons which are then transported into the lacteals within the intestinal villi and eventually empty into lymphatic duct. The lymphatic ducts ultimately release the absorbed substances into the blood stream. While the fatty acids are absorbed by the lymph duct, other materials are absorbed either actively or passively by the capillaries of the villi (Figure. 5.9). Water soluble vitamins are absorbed by simple diffusion or active transport. Transport of water depends upon the osmotic gradient.

Absorption of substances in the alimentary canal takes place in mouth, stomach, small intestine and large intestine. However maximum absorption takes place in the small intestine. Absorption of simple sugars, alcohol and medicines takes place in the stomach. Certain drugs are absorbed by blood capillaries in the lower side of the tongue and mucosa of mouth. Large intestine is also involved in absorption of more amounts of water, vitamins, some minerals and certain drugs.

Absorbed substances are transported through blood and lymph to the liver through the hepatic portal system. From the liver, nutrients are transported to all other regions of the body for utilization. All the body tissues utilize the absorbed




Figure 5.9 Process of Digestion and absorption

XI Std Zoology Chapter-5 EM.indd 82 30/12/2022 10:38:47