Proteins are source of amino acids required for growth and repair of body cells. They are stored in the body only to a certain extent large quantities are excreted as nitrogenous waste. The caloric value and physiological fuel value of one gram of protein are 5.65 Kcal and 4 Kcal respectively. According to ICMR (Indian Council of Medical Research and WHO (World Health Organization), the daily requirement of protein for an average Indian is 1gm per 1 kg body weight.

5.7 Nutritional and digestive disorders

Intestinal tract is more prone to bacterial, viral and parasitic worm infections. This infection may cause inflammation of the inner lining of colon called **colitis**. The most common symptoms of colitis are rectal bleeding, abdominal cramps, and diarrhoea.

Protein energy malnutrition: (PEM)

Growing children require more amount of protein for their growth and development. Protein deficient diet during early stage of children may lead to protein energy malnutrition such as **Marasmus and Kwashiorkor**. Symptoms are dry skin, pot-belly, oedema in the legs and face, stunted growth, changes in hair colour, weakness and irritability. Marasmus is an acute form of protein malnutrition. This condition is due to a diet with inadequate carbohydrate and protein. Such children are suffer from diarrhoea, body becomes lean and weak (emaciated) with reduced fat and muscle tissue with thin and folded skin.

Indigestion

It is a digestive disorder in which the food is not properly digested leading to a feeling of fullness of stomach. It may be due to inadequate enzyme secretion, anxiety, food poisoning, over eating, and spicy food.

Constipation

In this condition, the faeces are retained within the rectum because of irregular bowel movement due to poor intake of fibre in the diet and lack of physical activities.

Vomiting

It is reverse peristalsis. Harmful substances and contaminated food from stomach are ejected through the mouth. This action is controlled by the vomit centre located in the medulla oblongata. A feeling of nausea precedes vomiting.

Jaundice

It is the condition in which liver is affected and the defective liver fails to break down haemoglobin and to remove bile pigments from the blood. Deposition of these pigments changes the colour of eye and skin yellow. Sometimes, jaundice is caused due to hepatitis viral infections.

Liver Cirrhosis

Chronic disease of liver results in degeneration and destruction of liver cells resulting in abnormal blood vessel and bile duct leading to the formation of fibrosis. It is also called deserted liver or scarred liver. It is caused due to infection, consumption of poison, malnutrition and alcoholism.

Gall Stones

Any alteration in the composition of the bile can cause the formation of stones in the gall bladder. The stones are mostly formed of crystallized cholesterol in the bile. The gall stone causes obstruction in the cystic duct, hepatic duct and also hepato-pancreatic duct causing pain, jaundice and pancreatitis.

XI Std Zoology Chapter-5 EM.indd 86 30/12/2022 10:38:47