(synthesized in the liver) to angiotensin I. Angiotensin converting enzyme (ACE) converts angiotensin I to angiotensin II. Angiotensin II stimulates Na^+ reabsorption in the distal convoluted tubule by vasoconstriction of the blood vessels and increases the glomerular blood pressure. Angiotensin II acts at different sites such as heart, kidney, brain, adrenal cortex and blood vessels. It stimulates adrenal cortex to secrete aldosterone that causes reabsorption of Na^+, K^+ excretion and absorption of water from the distal convoluted tubule and collecting duct. This increases the glomerular blood pressure and glomerular filtration rate. This complex mechanism is generally known as **Renin - Angiotensin - Aldosterone System** (RAAS). Figure 8.9 shows the schematic representation of the various hormones in the regulation of body fluid concentration.

Atrial natriuretic factor

Excessive stretch of cardiac atrial cells cause an increase in blood flow to the atria of the heart and release Atrial Natriuretic Peptide or factor (ANF) travels to the kidney where it increases Na^+ excretion and increases the blood flow to the glomerulus, acting on the afferent glomerular arterioles as a vasodilator or on efferent arterioles as a vasoconstrictor. It decreases aldosterone release from the adrenal cortex and also decreases release of renin, thereby decreasing angiotensin II. ANF acts antagonistically to the renin- angiotensin system, aldosterone and vasopressin.

8.5 Micturition

The process of release of urine from the bladder is called micturition or urination. Urine formed by the nephrons is ultimately carried to the urinary bladder where it is stored till it receives a voluntary signal from the central nervous system. The stretch receptors present in the urinary bladder are stimulated when it gets filled with urine. Stretching of the urinary bladder stimulates the CNS via the sensory neurons of the parasympathetic nervous system and brings about contraction of the bladder. Simultaneously, somatic motor neurons induce the sphincters to close. Smooth muscles contracts resulting in the opening of the internal sphincters passively and relaxing the external sphincter. When the stimulatory and inhibitory controls exceed the threshold, the sphincter opens and the urine is expelled out.

An adult human on an average excretes 1 to 1.5 L of urine per day. The urine formed is a yellow coloured watery fluid which is slightly acidic in nature (pH 6.0), Changes in diet may cause pH to vary between 4.5 to 8.0 and has a characteristic odour. The yellow colour of the urine is due to the presence of a pigment, urochrome. On an average, 25-30 gms of

Hypotonic urine is formed when osmotic pressure of the body fluid is decreased due to water retention or solute loss when ADH secretion is lowered. If you drink large volume of water without eating anything salty, the total body fluid volume increases quickly and the osmolarity decreases. The kidneys increases the volume of urine excreted. The reverse happens when you eat salty food without drinking water.

XI Std Zoology Chapter-8 EM.indd 146 30/12/2022 13:48:08