(15) If $\Delta = \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix}$, then $\begin{vmatrix} ka & kb & kc \\ kx & ky & kz \\ kp & kq & kr \end{vmatrix}$ is

(1) Δ (2) $k\Delta$ (3) $3k\Delta$ (4) $k^3\Delta$

(16) A root of the equation $\begin{vmatrix} 3-x & -6 & 3 \\ -6 & 3-x & 3 \\ 3 & 3 & -6-x \end{vmatrix} = 0$ is

(1) 6 (2) 3 (3) 0 (4) – 6

(17) The value of the determinant of $A = \begin{bmatrix} 0 & a & -b \\ -a & 0 & c \\ b & -c & 0 \end{bmatrix}$ is

(1) $-2abc$ (2) abc (3) 0 (4) $a^2+b^2+c^2$

(18) If x_1, x_2, x_3 as well as y_1, y_2, y_3 are in geometric progression with the same common ratio, then the points (x_1, y_1), (x_2, y_2), (x_3, y_3) are

(1) vertices of an equilateral triangle

(2) vertices of a right angled triangle

(3) vertices of a right angled isosceles triangle

(4) collinear

(19) If $\lfloor . \rfloor$ denotes the greatest integer less than or equal to the real number under consideration and $-1 \le x < 0,\ 0 \le y < 1,\ 1 \le z < 2$, then the value of the determinant $\begin{vmatrix} \lfloor x \rfloor + 1 & \lfloor y \rfloor & \lfloor z \rfloor \\ \lfloor x \rfloor & \lfloor y \rfloor + 1 & \lfloor z \rfloor \\ \lfloor x \rfloor & \lfloor y \rfloor & \lfloor z \rfloor + 1 \end{vmatrix}$ is

(1) $\lfloor z \rfloor$ (2) $\lfloor y \rfloor$ (3) $\lfloor x \rfloor$ (4) $\lfloor x \rfloor + 1$

(20) If $a \neq b, b, c$ satisfy $\begin{vmatrix} a & 2b & 2c \\ 3 & b & c \\ 4 & a & b \end{vmatrix} = 0$, then $abc =$

(1) $a+b+c$ (2) 0 (3) b^3 (4) $ab+bc$

(21) If $A = \begin{vmatrix} -1 & 2 & 4 \\ 3 & 1 & 0 \\ -2 & 4 & 2 \end{vmatrix}$ and $B = \begin{vmatrix} -2 & 4 & 2 \\ 6 & 2 & 0 \\ -2 & 4 & 8 \end{vmatrix}$, then B is given by

(1) $B = 4A$ (2) $B = -4A$ (3) $B = -A$ (4) $B = 6A$

07_11th-Maths-Vol2-Unit7.indd 42 16-12-2022 10:59:56