and $y''' = (-1)(-2)(-3)x^{-4} = \dfrac{(-1)^3 3!}{x^4}$.

Example 10.33

Find f'' if $f(x) = x \cos x$.

Solution

We have, $f(x) = x \cos x$.

Now $f'(x) = -x \sin x + \cos x$, and

$f''(x) = -(x \cos x + \sin x) - \sin x$

$= -x \cos x - 2 \sin x$.

Example 10.34

Find y'' if $x^4 + y^4 = 16$.

Solution

We have $x^4 + y^4 = 16$.

Differentiating implicitly, $4x^3 + 4y^3 y' = 0$

Solving for y' gives

$$y' = -\frac{x^3}{y^3} .$$

To find y'' we differentiate this expression for y' using the quotient rule and remembering that y is a function of x.

$$y'' = \frac{d}{dx}\left(\frac{-x^3}{y^3}\right) = \frac{-\left[y^3 \dfrac{d}{dx}(x^3) - x^3 \dfrac{d}{dx}(y^3)\right]}{(y^3)^2}$$

$$= -\frac{\left[y^3 . 3x^2 - x^3 (3y^2 y')\right]}{y^6}$$

$$= -\frac{3x^2 y^3 - 3x^3 y^2 \left(-\dfrac{x^3}{y^3}\right)}{y^6}$$

$$= -\frac{3(x^2 y^4 + x^6)}{y^7} = \frac{-3x^2 [x^4 + y^4]}{y^7}$$

$$= \frac{-3x^2 (16)}{y^7} = \frac{-48x^2}{y^7} .$$

Example 10.35

Find the second order derivative if x and y are given by

$x = a \cos t$

$y = a \sin t$.

10_11th-Maths-Vol2-Unit10.indd 174 16-12-2022 11:05:31