www.tntextbooks.in

height h_1 to h_2, then the potential energy at h_1 is

$$U(h_1) = -mgR_e + mgh_1 \qquad (6.36)$$

and the potential energy at h_2 is

$$U(h_2) = -mgR_e + mgh_2 \qquad (6.37)$$

The potential energy difference between h_1 and h_2 is

$$U(h_2) - U(h_1) = mg(h_1 - h_2). \qquad (6.38)$$

The term mgR_e in equations (6.36) and (6.37) plays no role in the result. Hence in the equation (6.35) the first term can be omitted or taken to zero. Thus it can be stated that The gravitational potential energy stored in the particle of mass m at a height h from the surface of the Earth is $U = mgh$. On the surface of the Earth, $U = 0$, since h is zero.

It is to be noted that mgh is the work done on the particle when we take the mass m from the surface of the Earth to a height h. This work done is stored as a gravitational potential energy in the mass m. Even though mgh is gravitational potential energy of the system (Earth and mass m), we can take mgh as the gravitational potential energy of the mass m since Earth is stationary when the mass moves to height h.

6.2.5 Gravitational potential V(r)

It is explained in the previous sections that the gravitational field $\vec{E}$ depends only on the source mass which creates the field. It is a vector quantity. We can also define a scalar quantity called "gravitational potential" which depends only on the source mass.

The gravitational potential at a distance r due to a mass is defined as the amount of work required to bring unit mass from infinity to the distance r and it is denoted as V(r). In other words, the gravitational potential at distance r is equivalent to gravitational potential energy per unit mass at the same distance r. It is a scalar quantity and its unit is J kg^{-1}

We can determine gravitational potential from gravitational potential energy. Consider two masses m_1 and m_2 separated by a distance r which has gravitational potential energy $U(r)$ (Figure 6.15). The gravitational potential due to mass m_1 at a point P which is at a distance r from m_1 is obtained by making m_2 equal to unity (m_2 = 1kg). Thus the gravitational potential $V(r)$ due to mass m_1 at a distance r is

$$V(r) = -\frac{Gm_1}{r} \qquad (6.39)$$




Figure 6.15 Point mass placed at a distance

Gravitational field and gravitational force are vector quantities whereas the gravitational potential and gravitational potential energy are scalar quantities. The motion of particles can be easily analyzed using scalar quantities than vector quantities. Consider the example of a falling apple:

Figure 6.16 shows an apple which falls on Earth due to Earth's gravitational force. This can be explained using the concept of gravitational potential $V(r)$ as follows.