6. What is the difference between 56L and 56?

7. Determine which of the following are valid constant? And specify their type.

   (i) 0.5 (ii) 'Name' (iii) '\t' (iv) 27,822

8. Suppose x and y are two double type variable that you want add as integer and assign to an integer variable. Construct a C++ statement to do the above.

9. What will be the result of following if num=6 initially.

   (a) cout << num;

   (b) cout << (num==5);

10. Which of the following two statements are valid? Why? Also write their result.

    (i) int a; a = 3,014; (ii) int a; a=(3,014);

SECTION-C

Short Answers

1. What are arithmetic operators in C++? Differentiate unary and binary arithmetic operators. Give example for each of them.

2. How relational operators and logical operators are related to one another?

3. Evaluate the following C++ expressions where x, y, z are integers and m, n are floating point numbers. The value of x = 5, y = 4 and m=2.5;

   (i) n = x + y / x;

   (ii) z = m * x + y;

   (iii) z *= x * m + x;

Reference:

(1) Object Oriented Programming with C++ (4th Edition), Dr. E. Balagurusamy, Mc.Graw Hills.

(2) The Complete Reference C++ (Forth Edition), Herbert Schildt. Mc.Graw Hills.

(3) Computer Science with C++ (A text book of CBSE XI and XII), Sumita Arora, Dhanpat Rai & Co.

11th Std Computer Science_EM Chapter 9.indd 151 06-01-2023 15:08:46