its functions are protection, secretion and absorption. Ciliated forms line the trachea and the upper respiratory tract. The non ciliated forms, line the epididymis, large ducts of a glands and tracts of male urethra

Important epithelial tissue disorders:
Eczema, Psoriasis, Epithelial carcinoma and severe asthma

Glandular epithelium

Some of the cuboidal or columnar cells get specialized for secretion and are called **glandular epithelium** (Figure 3.3). They are mainly of two types: unicellular, consisting of isolated glandular cells (goblet cells of the **alimentary canal**), and multicellular, consisting of cluster of cells (**salivary gland**). On the basis of the mode of pouring of their secretions, glands are divided into two categories namely exocrine and endocrine glands. **Exocrine glands** secrete mucus, saliva, earwax, oil, milk, digestive enzymes and other cell products. These products are released through ducts or tubes. In contrast endocrine glands do not have ducts. Their secretions called hormones are secreted directly into the fluid bathing the gland. The exocrine glands are classified as unicellular and multicelluar glands. The multicelluar glands are further classified based on the structure as **simple** and **compound glands,** based on their secretory units as **tubular, alveolar (Acinus)** and **tubulo alveolar**. Based on the mode of secretion exocrine glands are classified as **merocrine, holocrine** and **apocrine**.





Figure 3.3 Glandular Epithelium



XI Std Zoology Chapter-3 EM.indd 40 30/12/2022 10:33:23