www.tntextbooks.in

The outline of the classification is given below:



1.5.5 Kingdom : Myceteae (Fungi)

Include achlorophyllous, saprophytic or parasitic organisms with Unicellular or multicellular (Mycelium) thallus surrounded by chitinous cell wall. Nutrition is absorptive except slime molds.Reproduction is through asexual and Sexual methods.

Division : I Gymnomycota

Nutrition Phagotrophic, members of this group lack cell wall. Example. *Dictyostelium*

Division :II Mastigomycota

Flagellate cells are present(Gamete/ Zoospore). Nutrition absorptive, mycelium coenocytic. Example : *Albugo*

Division : III Amastigomycota

Unicellular to multicellular forms are included. The mycelium is septate. Asexual reproduction occurs by budding, fragmentation, sporangiospores, conidia etc., Meiosis is zygotic. Example : *Peziza*

Recently, with the advent of molecular methods myxomycetes and oomycetes were reclassified and treated under chromista.

The salient features of some of the classes – Oomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Form class Deuteromycetes are discussed below.

Oomycetes

Coenocytic mycelium is present. The cell wall is made up of Glucan and Cellulose. Zoospore with one whiplash and one tinsel flagellum is present. Sexual reproduction is Oogamous. Example: *Albugo.*