www.tntextbooks.in

2.5 Pteridophytes

Seedless Vascular Cryptogams



From the previous chapter we are aware of the salient features of amphibious plants called Bryophytes. But there is a plant group called Pteridophytes which are considered as first true land plants. Further, they were the first plants to acquire vascular tissue namely xylem and phloem, hence called vascular cryptogams. Club moss, Horsetails, quill worts, water ferns and Tree ferns belong to this group. This chapter deals with the characteristic features of Pteridophytes.

Pteridophytes are the vascular cryptogams and were abundant in the Devonian period of Palaeozoic era (400 million years ago). These plants are mostly small, herbaceous and grow well in moist, cool and shady places where water is available. The photographs for some pteridophytes are given in Figure 2.24.

2.5.1 General characteristic features of Pteridophytes:

- Plant body is sporophyte (2n) and it is the dominant phase. It is differentiated into root, stem and leaves.
- Roots are adventitious.
- Stem shows monopodial or dichotomous branching.
- Leaves may be microphyllous or megaphyllous.
- Stele is protostele but in some forms siphonostele is present (*Marsilea*)
- Tracheids are the major water conducting elements but in *Selaginella* vessels are found.
- Sporangia, spore bearing bag like structures are borne on special leaves called sporophyll. The sporophylls gets organized to form cone or strobilus. Example: *Selaginella*, *Equisetum* .
- They may be **homosporous** (produce one type of spores-*Lycopodium*) or **Heterosporous** (produce two types of spores-*Selaginella)*. Heterospory is the origin for seed habit.
- Development of sporangia may be **eusporangiate** (development of sporangium from group of initials) or **leptosporangiate** (development of sporangium from single initial).
- Spore mother cells undergo meiosis and produce spores (n).
- Spore germinates to produce haploid, multicellular green, cordate shaped independent gametophytes called prothallus.
- Fragmentation, Resting buds, root tubers and adventitious buds help in Vegetative reproduction.
- Sexual reproduction is Oogamous. Sex organs, namely antheridium and archegonium are produced on the prothallus.
- Antheridium produces spirally coiled and multiflagellate antherozoids.
- Archegonium is flask shaped with broad venter and elongated narrow neck. The venter possesses egg or ovum and neck contain neck canal cells.