www.tntextbooks.in

Radicle doesn't persist and fibrous root is present.

Flowers trimerous.

Monocolpate (1 furrow) Pollen is present.

Anatomical features

- Vascular bundles are scattered in the stem
- Vascular bundles are closed (Cambium absent).
- Secondary growth is absent.

Current Angiosperm Phylogeny Group (APG) System of classification doesn't recognize dicots as a monophyletic group. Plants that are traditionally classified under dicots are dispersed in several clades such as early Magnolids and Eudicots.

Summary

Plant Kingdom includes Algae, Bryophytes, Pteridophytes, Gymnosperms and Angiosperms

The life cycle in plants fall under three types 1. Haplontic,2. Diplontic and 3.Haplodiplontic

Algae are autotrophic, chlorophyll bearing organisms. The Plant body is not differentiated into root like, stem like or leaf like structures. A wide range of thallus organization is found in algae. They reproduce vegetatively through fragmentation, tuber and akinete formation. Zoospores , autospores and hypnospores are produced during asexual reproduction .and Sexual reproduction occurs through isogamy, anisogamy and oogamy.

Oedogonium is a fresh water, filamentous, multicellular alga. The presence of cap cell is the prominent characteristic feature in addition reticulate chloroplast is present. Asexual reproduction takes place through Zoospores. The sexual reproduction is Oogamous.

Chara is a fresh water alga and is popularly called "Stone worts". The plant body is multicellular, macroscopic and is differentiated into main axis and rhizoids. Sexual reproduction is Oogamous.

Bryophytes are simplest land plants. They are called amphibians of plant kingdom or nonvascular cryptogams. The plant body is gametophyte. The sporophyte depends upon gametophyte. Conducting tissues like xylem and phloem is absent. Vegetative reproduction takes place through fragmentation, formation of adventitious bud and Gemmae. Sexual reproduction is Oogamous. Water is essential for fertilization.

Marchantia belongs to the class Hepaticopsida. The thallus is dorsiventral and is attached to the substratum by means of rhizoids. The internal structure of the thallus reveals the presence of photosynthetic region and a storage region. Vegetative reproduction takes place through fragmentation and formation of Gemmae. The sexual reproduction is Oogamous. Sporophyte bears spores. Alternation of generation is present.

Funaria belongs to the class Bryopsida. The gametophyte is differentiated into leaf-like, stem-like structures with rhizoids. Gemmae, Protonema and bulbils help in asexual reproduction. Sexual reproduction is Oogamous. Alternation of generation is present.

Pteridophytes are also called vascular cryptogams. The plant body is sporophyte and is long lived ,which is differentiated into root, stem and leaves. They may be homosporous or heterosporous.