www.tntextbooks.in

3. Sucking or Haustorial roots

These roots are found in parasitic plants. Parasites develop adventitious roots from stem which penetrate into the tissue of the host plant and suck nutrients.

Example: *Cuscuta* (dodder), *Cassytha*, *Orobanche* (broomrape), *Viscum* (mistletoe), *Dendrophthoe*.

4. Photosynthetic or assimilatory roots

Roots of some climbing or epiphytic plants develop chlorophyll and turn green which help in photosynthesis. Example: *Tinospora*, *Trapa natans* (water chestnut), *Taeniophyllum*.

3.6 Shoot system

The plumule of the embryo of a germinating seed grows into stem. The epicotyl elongates after embryo growth into the axis (the stem) that bears leaves from its tip, which contain the actively dividing cells of the shoot called **apical meristem.** Further cell divisions and growth result in the formation of mass of tissue called **a leaf primordium.** The point from which the leaf arises is called **node**. The region between two adjacent nodes is called **internode.**

I. Characteristic features of the stem

1. The stem is usually the aerial portion of the plant
2. It is positively phototropic and negatively geotropic
3. It has nodes and internodes.
4. Stem bears vegetative bud for vegetative growth of the plant, and floral buds for reproduction, and ends in a terminal bud.
5. The young stem is green and thus carries out photosynthesis.
6. During reproductive growth stem bears flowers and fruits.
7. Branches arise exogenously
8. Some stems bears multicellular hairs of different kinds.

II. Functions of the stem

Primary functions

1. Provides support and bears leaves, flowers and fruits.
2. It transports water and mineral nutrients to the other parts from the root.
3. It transports food prepared by leaves to other parts of the plant body.



Vanda *Bryophyllum* *Cuscuta* *Tinospora*

Figure 3.7: Adventitious Root Modification for Vital Functions