www.tntextbooks.in

f. Panicle: A branched raceme is called **panicle**. Example: *Mangifera,* neem, *Delonix regia*. It is also called **Compound raceme** or **raceme of racemes**.

2. Main axis shortened:

Inflorescence with reduced growth of central axis. There are two types namely corymb and umbel.

a. Corymb: An inflorescence with shorter pedicellate flowers at the top and longer pedicellate flowers at the bottom. All flowers appear at the same level to form convex or flat topped racemose inflorescence. Example: *Caesalpinia*. **Compound corymb:** A branched corymb is called **compound corymb**. Example: Cauliflower.




Figure 4.4: (m) Corymb diagrammatic




Figure 4.4: (n) Compound corymb diagrammatic



Figure 4.4: (o) Corymb



Figure 4.4: (p) Compound corymb




Figure 4.4: (q) Umbel diagrammatic




Figure 4.4: (r) Compound umbel diagrammatic

b. Umbel: An inflorescence with indeterminate central axis and pedicellate flowers arise from a common point of peduncle at the apex. Example: *Allium cepa*, *Centella asiatica*, *Memecylon umbellatum*. **Compound umbel**: It is a branched umbel. Each smaller unit is called **umbellule**. Example: *Daucas carota*, *Coriandrum sativum, Memecylon edule*.



Figure 4.4: (s) Umbel



Figure 4.4: (t) Compound umbel

3. Main axis flattened:

The main axis of inflorescence is mostly flattened (convex or concave) or globose. A **head** or **capitulum** is a determinate or indeterminate, group of sessile or sub sessile flowers arising on a receptacle, often subtended by an involucre.

a. Head: A head is a characteristic inflorescence of Asteraceae and is also found in some members of Rubiaceae.



Figure 4.4: (u) *Neolamarkia cadamba* head

Example: *Neolamarkia cadamba*, *Mitragyna parvifolia* and in some members of Fabaceae-Mimosoideae. Example: