www.tntextbooks.in




Figure 4.6: (b) diagrammatic,
(c) Monochasial Helicoid

b. Scorpioid: Axis develops on alternate sides and often becomes a coil structure. Example: *Heliotropium*.



Figure 4.6: (d) diagrammatic,
(e) Monochasial Scorpioid

3. Simple dichasium (Biparous): A central axis ends in a terminal flower; further growth is produced by two lateral buds. Each cymose unit consists of three flowers of which central one is old one. This is **true cyme**. Example: *Jasminum*.



Figure 4.6: (f) diagrammatic,
(g) Simple dichasium

4. **Compound dichasium**: It has many flowers. A terminal old flower develops lateral simple dichasial cymes on both sides. Each compound dichasium consists of seven flowers. Example: *Clerodendron*.

A small,simple dichasium is called **cymule**




Figure 4.6: (h) diagrammatic,
(i) Compound dichasium

5. Polychasial Cyme (multiparous): The central axis ends with a flower. The lateral axes branches repeatedly. Example: *Nerium*



Figure 4.6: (j) diagrammatic,
(k) Polychasial cyme



Sympodial Cyme: In monochasial cyme, successive axes at first develop in a zigzag manner and later it develops into a straight pseudo axis. Example: *Solanum americanum*.

III. Mixed Inflorescence




Figure 4.7: Mixed and special inflorescence