www.tntextbooks.in

1. Uniseriate: It is a single whorl of accessory(non-essential) floral part. It is less common.Example: *Sterculia.*

2. Biseriate (dicyclic): It is with two whorls of accessory floral parts. (outer=lower,inner=upper)It is the most common type of perianth cycly. Example: *Hibiscus.*

3. Multiseriate: (**triseriate,tetraseriate**) More than two whorls of non–essential floral parts.Example: *Chrysanthemum.*

4. Dichlamydeous: A flower is composed of distinct outer calyx and inner corolla.

5. Homochlamydeous: Perianth is undifferentiated into calyx and corolla and composed of similar parts called **tepals**. Most monocots have a homochlamydeous perianth.

6. Achlamydeous: Perianth is absent altogether.Flowers without petals are called **apetalous** and flowers without sepals are called **asepalous.**



Figure 4.15: (a)
Uniseriate



Figure 4.15: (b)
Biseriate



Figure 4.15: (c)
Multiseriate



Figure 4.16: (a)
Dichlamydeous



Figure 4.16: (b)
Homochlamydeous

4.3.3 Merosity

Number of floral parts per whorl is called **merosity**. Perianth merosity is the number of perianth parts per whorl.

1. Isomerous: Presence of same number of perianth parts in different whorls of a flower. (five sepals, five petals). Example: *Hibiscus*.

2. Anisomerous: Each whorl of flower contains different number of members. Example: *Annona*.

3. Bimerous: Floral parts in two or multiples of two. Example: *Ixora*

4. Trimerous: Floral parts in three or multiples of three. Example: *Allium*, Monocots.

5. Tetramerous: Floral parts in four or multiples of four. Example: *Brassica juncea.*

6. Pentamerous: Floral parts in five or multiples of five. Example: *Hibiscus*, Dicots.

4.3.4 Calyx

Calyx protects the flower in bud stage. Outermost whorl of flower is calyx. Unit of calyx is sepal. Normally green in colour.