IV. Sympetalous Zygomorphic

1. Bilabiate:

Corolla with two lips. Example: *Ocimum,Leucas,Adhatoda*.

Tubular corolla with a single strap-shaped limb. Example: Ray floret of *Helianthus*

2. Personate:

Corolla made up of two lips with the upper arched and the lower protruding into the corolla throat. Example: *Antirrhinum,Linaria*.

3. Ligulate:

Tubular corolla with a single strap-shaped limb. Example: Ray floret of *Helianthus*.



Figure 4.21: (i) Bilabiate



Figure 4.21: (j) Personate

4.3.6 Perianth

Can you recall the term homochlamydeous? undifferentiated calyx and corolla in a flower is called **perianth**. Each member is called **tepal**. If the tepals are distinct they are called **Apotepalous** (Polyphyllous). Example: *Allium sativum*. Fused tepals are called **Syntepalous**. (Gamophyllous). Example: *Allium cepa*.

C0Xcnxcvg< Margins of sepals or petals do not overlap but just touch each other.
Example: Calyx in members of Malvaceae, *Calotropis, Annona.*

B. **Vy kuvgf " qt" eqpxqnvwg" qt" eqpvqtvgf <**
One margin of each petal or sepal overlapping on the other petal
Example: Petals of chinarose

Cguvkxcvlqp"
Arrangement of sepals and petals in the flower bud.

C. Imbricate: Sepals and petals irregularly overlap on each other; one member of the whorl is exterior, one interior and rest of the three having one margin exterior and the other interior.
Example: *Cassia, Delonix*
There are 3 types.
1.Ascendingly imbricate.
2.Quincuncial.
3.Vexillary.

D. Quincuncial: It is a type of imbricate aestivation in which two petals are external and two internal and one petal with one margin internal and the other margin external.
Example:Guava,calyx of *Ipomoea, Catharanthus*.

E. Vexillary: Large posterior petals both margins overlap lateral petals.
Lateral petals other margin overlaps anterior petals
Example: Pea,bean.