www.tntextbooks.in




Figure 4.40: Simple fleshy fruits

superior ovary. The fruit wall is differentiated into leathery epicarp with oil glands, a middle fibrous mesocarp. The endocarp forms distinct chambers, containing juicy hairs. Example: Orange, Lemon.

e) Pome: It develops from multicarpellary, syncarpous, inferior ovary. The receptacle also develops along with the ovary and becomes fleshy, enclosing the true fruit. In pome the epicarp is thin skin like and endocarp is cartilagenous. Example: Apple, Pear.

f) Balausta: A fleshy indehiscent fruit developing from multicarpellary, multilocular inferior ovary whose pericarp is tough and leathery. Seeds are attached irregularly with testa being the edible portion. Example: Pomegranate.

B. Dry Fruit

They develops from single ovary where the pericarp is dry and not differentiated into epicarp, mesocarp and endocarp. It is further subdivided into three types.

1) Dry dehiscent fruit

Pericarp is dry and splits open along the sutures to liberate seeds. They can be classified into following types.

a) Follicle: Fruit develops from monocarpellary, superior ovary and dehisces along one suture. Example: *Calotropis.*

b) Legume or pod: Fruit develops from monocarpellary, superior ovary and dehisces through both dorsal and ventral sutures. Example: *Pisum.*

c) Siliqua: Fruit develops from bicarpellary, syncarpous, superior ovary initially one chambered but subsequently becomes two chambered due to the formation of false septum (**replum**). The fruit dehisces along two suture. Example: *Brassica.*