www.tntextbooks.in



Follicle (*Calotropis*)



Legume (*Pisum*)



Siliqua (*Brassica*)



Silicula (*Capsella*)



Loculicidal
(Lady's finger)



Septifragal (*Datura*)

Figure 4.41: Dry dehiscent fruit

d) Silicula: Fruit similar to siliqua but shorter and broader. Example: *Capsella, Lepidium, Alyssum.*

e) Capsule: Fruit develops from multicarpellary, syncarpous, superior ovary. Based on the dehiscence pattern they are sub divided into.

i) Septicidal: Capsule splitting along septa and valves remaining attached to septa. Example: *Linum, Aristolochia.*

ii) Loculicidal: Capsule splitting along locules and values remaining attached to septa. Example: Lady's finger.

iii) Septifragal: Capsule splitting so that valves fall off leaving seeds attached to the central axis. Example: *Datura.*

iv) Poricidal: Dehiscence through terminal pores. Example: *Papaver.*

v) Denticidal: Capsule opening at top exposing a number of teeth. Example: *Primula, Cerastium.*

vi) Circumscissile: (pyxidium) Dehisces transversely so that top comes off as a lid or operculum. Example: *Anagallis arvensis, Portulaca, Operculina.*

2) Dry indehiscent fruit

Dry fruit which does not split open at maturity. It is subdivided into.

a) Achene: Single seeded dry fruit developing from single carpel with superior ovary. Achenes commonly develop from apocarpous pistil, Fruit wall



Achene (*Clematis*)



Cypsela (*Tridax*)



Caryopsis (*Oryza*)



Nut (*Anacardium*)



Samara (*Acer*)



Utricle
(*Chenopodium*)

Figure 4.42: Dry indehiscent fruit