www.tntextbooks.in

is free from seed coat. Example: *Clematis, Delphinium,* Strawberry.

b) Cypsela: Single seeded dry fruit, develops from bicarpellary, syncarpous, inferior ovary with reduced scales, hairy or feathery calyx lobes. Example: *Tridax, Helianthus.*

c) Caryopsis: It is a one seeded fruit which develops from a monocarpellary, superior ovary. Pericarp is inseparably fused with seed. Example: *Oryza, Triticum.*

d) Nut: They develop from mulicarpellary, syncarpous, superior ovary with hard, woody or bony pericap. It is a one seeded fruit. Example: *Quercus, Anacardium.*

e) Samara: A dry indehiscent, one seeded fruit in which the pericarp devlops into thin winged structure around the fruit. Example: *Acer*, *Pterocarpus.*

f) Utricle: They develops from bicarpellary, unilocular, syncarpus, superior ovary with pericarp loosely enclosing the seeds. Example: *Chenopodium.*

3) Schizocarpic Fruit

This fruit type is intermediate between dehiscent and indehiscent fruit. The fruit instead of dehiscing rather splits into number of segments, each containing one or more seeds. They are of following types.

a) Cremocarp: Fruit develops from bicarpellary, syncarpous, inferior ovary and splitting into two one seeded segments known as **mericarps**. Example: Coriander, Carrot.

b) Carcerulus: Fruit develops from bicarpellary, syncarpous, superior ovary and splitting into four one seeded segments known as **nutlets**. Example: *Leucas, Ocimum, Abutilon*



Cremocarp (Coriander)



Carcerulus (*Abutilon*)



Lomentum (Mimosa)



Regma (Castor)

Figure 4.43: Schizocarpic Fruit

c) Lomentum: The fruit is derived from monocarpellary, unilocular ovary. A leguminous fruit, constricted between the seeds to form a number of one seeded compartments that separate at maturity. Example: *Desmodium, Mimosa*

d) Regma: They develop from tricarpellary, syncarpous, superior, trilocular ovary and splits into one-seeded cocci which remain attached to carpophore. Example: *Ricinus, Geranium*

Aggregate Fruits

Aggregate fruits develop from a single flower having an apocarpous pistil. each of the free carpel is develops into a simple fruitlet. A collection of simple fruitlets makes an **aggregate fruit**. An individual ovary develops into a drupe, achene, follicle or berry. An aggregate of these fruits borne by a single flower is known as