www.tntextbooks.in

Gynoecium is female part of flower. Ovary, style and stigma are parts of pistil. According to number of carpels it is divided into monocarpellary, bicarpellary etc. It may be apocarpous or syncarpous. Locule number may be one to many. The ovary is superior or inferior or semi inferior. Mode of distribution of placenta inside the ovary is placentation. Construction of floral diagram and floral formula for given flower with some examples.

Fruits are the products of pollination and fertilization. Fruit developed from single ovary of flower is called **simple fruit**. Simple fruits are two types based on the fruit wall as simple fleshy and simple dry. An intermediate between dehiscent and indehiscent fruit is called schizocarpic fruit. The simple fruits could be fleshy or dry which could again be dehiscent or indehiscent. Fruits that are developed from multicarpellary, apocarpus pistil is called aggregate. Multiple or composite fruit develops from the flowers of the complete inflorescence. Seed is a ripened ovule which contains the embryo or the miniature of plant body. Seeds with one cotyledon are monocotyledonous and with two cotyledons are dicotyledonous.

Evalution

1. Vexillary aestivation is characteristic of the family
   a. Fabaceae
   b. Asteraceae
   c. Solanaceae d. Brassicaceae
2. Gynoecium with united carples is termed as
   a. Apocarpous b. Multicarpellary
   c. Syncarpous d. None of the above
3. Aggregate fruit develops from
   a. Multicarpellary, apocarpous ovary
   b. Multicarpellary, syncarpous ovary
   c. Multicarpellary ovary
   d. Whole inflorescence
4. In an inflorescence where flowers are borne laterally in an acropetal succession the position of the youngest floral bud shall be
   a. Proximal b. Distal
   c. Intercalary d. Anywhere
5. A true fruit is the one where
   a. Only ovary of the flower develops into fruit
   b. Ovary and calyx of the flower develops into fruit
   c. Overy, calyx and thalamus of the flower develops into fruit
   d. All floral whorls of the flower develops into fruit
6. Find out the floral formula for a bisexual flower with bract, regular, pentamerous, distinct calyx and corolla, superior ovary without bracteole.
7. Give the technical terms for the following: -
   a. A sterile stamen
   b. Stamens are united in one bunch
   c. Stamens are attached to the petals
8. Explain the different types of placentation with example.
9. Differenciate between aggregate fruit with multiple fruit.
10. Explain the different types of fleshy fruit with suitable example.