www.tntextbooks.in

the form of a tree called **phylogenetic tree.** Different molecular markers like allozymes, mitochondrial DNA, micro satellites, RFLP (Restriction Fragment Length Polymorphism), RAPD (Random amplified polymorphic DNA), AFLPs (Amplified Fragment Length Polymorphism), single nucleotide polymorphism- SNP, microchips or arrays are used in analysis.

Uses of molecular taxonomy

1. Molecular taxonomy helps in establishing the relationship of different plant groups at DNA level.
2. It unlocks the treasure chest of information on evolutionary history of organisms.

RFLP (Restriction Fragment Length Polymorphism)

RFLPs is a molecular method of genetic analysis that allows identification of taxa based on unique patterns of restriction sites in specific regions of DNA. It refers to differences between taxa in restriction sites and therefore the lengths of fragments of DNA following cleavage with restriction enzymes.

Amplified Fragment Length Polymorphism (AFLP)

This method is similar to that of identifying RFLPs in that a restriction enzyme is used to cut DNA into numerous smaller pieces, each of which terminates in a characteristic nucleotide sequence due to the action of restriction enzymes.

AFLP is largely used for population genetics studies, but has been used in studies of closely related species and even in some cases, for higher level cladistic analysis.

Random Amplified Polymorphic DNA (RAPD)

It is a method to identify genetic markers using a randomly synthesized primer that will anneal (recombine (DNA) in the double stranded form) to complementary regions located in various locations of isolated DNA. If another complementary site is present on the opposing DNA strand at a distance that is not too great (within the limits of PCR) then the reaction will amplify this region of DNA.

RAPDs like microsatellites may often be used for genetic studies within species but may also be successfully employed in phylogenetic studies to address relationships within a species or between closely related species. However RAPD analysis has the major disadvantage that results are difficult to replicate and in that the homology of similar bands in different taxa may be nuclear.

Significance of Molecular Taxonomy

1. It helps to identify a very large number of species of plants and animals by the use of conserved molecular sequences.
2. Using DNA data evolutionary patterns of biodiversity are now investigated.
3. DNA taxonomy plays a vital role in phytogeography, which ultimately helps in genome mapping and biodiversity conservation.
4. DNA- based molecular markers used for designing DNA based molecular probes, have also been developed under the branch of molecular systematics.

5.11.6 DNA Barcoding

Have you seen how scanners are used in supermarkets to distinguish the **Universal Product Code** (UPC)? In the same way we